

COGNITION

Clinical Research

Abiru, M., Mihara, Y., Kikuchi, Y., et al. 2007. The effects of Neurologic music therapy on hemispatial neglect in a hemiparetic stroke patient. A case study. *Neurological Medicine*, 67(1), 88-94.

Barrow, I.M, Collins, J.N., & Britt, L.D. 2006. The influence of an auditory distraction on rapid naming after a mild traumatic brain injury: a longitudinal study. *Journal of Trauma-Injury Infection & Critical Care*, 61, 1142-1149.

Baur, B., Uttner, I., Ilmberger, J., Fesl, G., & Mai, N. 2000. Music memory provides access to verbal knowledge in a patient with global amnesia. *Neurocase*, 6, 415-421.

Ben-Pazi, H., Gross-Tsur, V., Bergman, H., Shalev, R.S., (2003). Abnormal rhythmic motor response in children with attention-deficit-hyperactivity disorder. *Developmental Medicine & Child Neurology*. Vol. 45 pp.743-745.

Berel, M., Diller, L., & Orgel, M. 1971. Music as a facilitator for visual motor sequencing tasks in children with cerebral palsy. *Developmental Medicine and Child Neurology*, 13, 335-342.

Bettison, S. 1996. The long-term effects of auditory training on children with autism. *Journal of Autism and Developmental Disorders*, 26 (3), 361-375.

Bialystok, E. & Depape, A.M. 2009. Musical expertise, bilingualism, and executive functioning. *Journal of Experimental Psychology: Human Perception & Performance*, 35, 565-574.

Boso, M., Emanuele, E., Minazzi, V., Abbamonte, M., & Politi, P. 2007. Effect of long-term interactive music therapy on behavior profile and musical skills in young adults with severe autism. *The Journal of Alternative and Complimentary Medicine*, 13 (7), 709-712.

Brownell, M. 2002. Musically adapted social stories to modify behaviors in students with autism: Four case studies. *Journal of Music Therapy*, 39 (2), 117-144.

Bugos, J.A., Perlstein, W.M., McCrae, C.S., Brophy, T.S., Bedenbaugh, P.H. 2007. Individualized piano instruction enhances executive functioning and working memory in older adults. *Aging and Mental Health*, 11 (4), 464-71.

Carlomagno, S., Van Eeckhout, P., Blasi, V., Belin, P., Samson, Y., & Deloche, G. 1997. The impact of functional neuroimaging methods on the development of a theory for cognitive remediation. *Neuropsychological Rehabilitation*, 7, 311-326.

Carr, C. et al. (2012). Group music therapy for patients with persistent post-traumatic stress disorder—An exploratory randomized controlled trial with mixed methods evaluation. *Psychology and Psychotherapy: Theory, Research and Practice*, 85(2), 179-202.

Cavaco S, Feinstein JS, van Twillert H, Tranel D. 2012. Musical memory in a patient with severe anterograde amnesia. *J Clin Exp Neuropsychol*. 2012 Oct 4. [Epub ahead of print]

Ceccato, E., Caneva, P., Lamonaca, D. 2006. Music therapy and cognitive rehabilitation in schizophrenic patients: A controlled study. *Nordic Journal of Music Therapy*, 15(2), 111-120.

Chan, A.S., Ho, Y.C., & Cheung, M.C. 1998. Music training improves verbal memory. *Nature*, 396, 128.

Claussen, D. & Thaut, M.H. 1997. Music as a mnemonic device for children with learning disabilities. *Canadian Journal of Music Therapy*, 5, 55-66.

Cook, M. et al. (2010). A randomized controlled trial exploring the effect of music on agitated behaviors and anxiety in older people with dementia. *Aging and Mental Health*, 14(8), 905-916.

Cowles, A., Beatty, W.W., Nixon, S.J., Lutz, L.J., Paulk, J., Paulk, K., & Ross, E.D. 2003. Musical skill in dementia: A violinist presumed to have Alzheimer's disease learns to play a new song. *Neurocase*, 9, 493-503.

Cuddy, L., Duffin, J. 2005. Music, memory, and Alzheimer's disease: is music recognition spared in dementia, and how can it be assessed? *Medical Hypotheses*, 64, 229-235.

Davis, W.B., Gfeller, K., & Thaut, M.H. 1992 (1998, 2nd ed.). *An Introduction to Music Therapy: Theory and Practice*. Dubuque, IA: Wm. C. Brown Publishers. (Japanese Language Edition, 1997. Sapporo, Japan: Ichibaku Shuppansya Publishing.) (Spanish Language Edition, 2000) (Korean Language Edition, 2002)

Del Olmo, Aria P., Furio, M., Pozo, M., & Cudeiro, J. 2006. Evaluation of the effect of training using auditory stimulation on rhythmic movement in Parkinsonian patients--a combined motor and [18F]-FDG PET study. *Parkinsonism & Related Disorders*, 12(3), 155-64.

Dellacherie D, Bigand E, Molin P, Baulac M, Samson S. 2011. Multidimensional scaling of emotional responses to music in patients with temporal lobe resection. *Cortex*. 47(9),1107-15.

[Depape AM](#), [Hall GB](#), [Tillmann B](#), [Trainor LJ](#). 2012. Auditory processing in high-functioning adolescents with autism spectrum disorder. *PLoS One*. 7(9), e44084.

Depperschmidt, K. A.W. 1992. Musical mnemonics as an aid to memory in patients with dementia of Alzheimer's type. Unpublished master thesis, Colorado State University.

Dozza, M., Chiari, L., Hlavacka, F., Cappello, A., & Horak, F.B. 2006. Effects of linear versus sigmoid coding of visual or audio biofeedback for the control of upright stance. *IEEE Transactions on Neural Systems & Rehabilitation Engineering*, 14(4), 505-12.

Edgerton, C. 1994. The effect of improvisational music therapy on the communicative behavior of autistic children. *Journal of Music Therapy*, 31 (1), 31-62.

El Haj, M. et al. (2012). Music enhances autobiographical memory in mild Alzheimer's disease. *Education Gerontology*, 38(1), 30-41.

[El Haj M, Fasotti L, Allain P. 2012. The involuntary nature of music-evoked autobiographical memories in Alzheimer's disease. *Conscious Cogn.* 21\(1\),238-46.](#)

Erkkilä J, Gold C, Fachner J, Ala-Ruona E, Punkanen M, Vanhala M. (2008)The effect of improvisational music therapy on the treatment of depression: protocol for a randomised controlled trial. *BMC Psychiatry*. 8,50.

Field, T., Martinez, A., Nawrocki, T., Pickens, J., Fox, N.A., & Schanberg, S. 1998. Music shifts frontal EEG in depressed adolescents. *Adolescence*, 33, 109-116.

[Finnigan E, Starr E. 2010. Increasing social responsiveness in a child with autism. A comparison of music and non-music interventions. *Autism.* 14\(4\),321-48.](#)

[Ford JH, Addis DR, Giovanello KS. 2011. Differential neural activity during search of specific and general autobiographical memories elicited by musical cues. *Neuropsychologia.* 49\(9\),2514-26.](#)

Foster, N. & Valentine, E. 2001. The effect of auditory stimulation on autobiographical recall in dementia. *Experimental Aging Research*, 27, 215-228.

Forsblom, A., Laitinen, S., Sarkamo, T. & Tervaniemi, M. 2009. Therapeutic role of music listening in stroke rehabilitation. *Annals of the New York Academy of Sciences*, 1169, 426-430.

Fujioka, T., Ross, B., Kakigi, R., Pantey C., Trainor, LJ. 2006. One year of musical training affects development of auditory cortical-evoked fields in young children. *Brain*, 129, 2593-608.

Fukui H, Arai A, Toyoshima K. 2012. Efficacy of music therapy in treatment for the patients with Alzheimer's disease. *International Journal of Alzheimer's Disease*. Epub before print: Sept 26 2012.

Gaston, E.T. (Ed.). 1968. *Music in Therapy*. New York, NY: MacMillan Co.

Gerard, C. & Drake, C. 1990. The inability of young children to reproduce intensity differences in musical rhythms. *Perception and Psychophysics*, 48, 91-101.

Gfeller, K.E. 1983. Musical mnemonics as an aid to retention with normal and learning disabled students. *Journal of Music Therapy*, 20, 179-189.

Gfeller, K., Woodworth, G., Robin, D.A., Witt, S., & Knutson, J.F. 1997. Perception of rhythmic and sequential pitch patterns by normally hearing adults and adult cochlear implant users. *Ear and Hearing*, 18, 252-260.

Giacobbe, G.A. 1972. Rhythm builds order in brain-damaged children. *Music Educators Journal*, 58, 40-43.

Goerlich KS, Witteman J, Aleman A, Martens S. 2011. Hearing feelings: affective categorization of music and speech in alexithymia, an ERP study. *PLoS One*. 6(5),e19501.

Graham, Janet. 2004. Communication with the uncommunicative: Music therapy with pre-verbal adults. *British Journal of Learning Disabilities*, 32(1), 24-29.

Gregory, D. 2003. Music listening for maintaining attention of older adults with cognitive impairments. *Journal of Music Therapy*, 39 (4), 244-264.

Gronwall, D. 1977. Paced Auditory Serial-Addition Task: A measure of recovery from concussion. *Perceptual and Motor Skills*, 44, 367-373.

Guetin, S., Soua, B., Voiriot, G., Picot, M.C., & Herisson, C. 2009. The effect of music therapy on mood and anxiety-depression: an observational study in institutionalized patients with traumatic brain injury. *Annals of Physical & Rehabilitation Medicine*, 52, 30-40.

Hammar, L. et al. (2011). Communicating through caregiver singing during morning care situations in dementia care. *Scandinavian Journal of Caring Sciences*, 25(1), 160-168.

[Han P, Kwan M, Chen D, Yusoff SZ, Chionh HL, Goh J, Yap P. 2011. A controlled naturalistic study on a weekly music therapy and activity program on disruptive and depressive behaviors in dementia. *Dement Geriatr Cogn Disord*. 30\(6\),540-6.](#)

Haslam, C. & Cook, M. 2002. Striking a chord with amnesic patient's: Evidence that song facilitates memory. *Neurocase*, 8, 453-465.

Heaton, P., Hermelin, B., & Pring, L. 1988. Autism and pitch processing: A precursor for savant musical ability? *Music Perception*, 15, 291-305.

Hermelin, B., O'Connor, N., & Lee, S. 1987. Musical inventiveness of five idiots-savants. *Psychological Medicine*, 17 (3), 685-694.

Hermelin, B., O'Connor, N., Lee, S., & Treffert, D. 1989. Intelligence and musical improvisation. *Psychological Medicine*, 19 (2), 447-457.

Hiller, P.U. 1989. Song story: A potent tool for cognitive and affective relearning in head injury. *Cognitive Rehabilitation*, 7, 20-23.

Ho, Y., Cheung, M.C., & Chan, A.S. 2003. Music training improves verbal but not visual memory: Cross-sectional longitudinal explorations in children. *Neuropsychology*, 17 (3), 439-450.

Hommel, M., Peres, B., Pollak, P., Memin, B., Besson, G., Gaio, J.M., & Perret, J. 1990. Effects of passive tactile and auditory stimuli on left visual neglect. *Archives of Neurology*, 47, 573-576.

Hughes, J.R., Fino, J.J., & Melyn, M.A. 1999. Is there a chronic change of the "Mozart effect" on epileptiform activity? A case study. *Clinical Electroencephalography*, 30, 44-45.

Hummelsheim, H. 1999. Rationales for improving motor function. *Current Opinion in Neurology*, 12, 697-701.

Hurt-Thaut, C.P. & Johnson, S.B. 2003. Neurologic music therapy with children: Scientific foundations and clinical applications. In S.L. Robb (Ed.), *Music Therapy in Pediatric Healthcare: Research and Evidence-Based Practice* (pp. 81-100). Silver Spring, MD: American Music Therapy Association, Inc.

Irish, M., Cunningham, C.J., Walsh, J.B., Coakley, D., Lawlor, B.A., Robertson I.H., Coen, R.F. 2006. Investigating the enhancing effect of music on autobiographical memory in mild Alzheimer's disease. *Dementia & Geriatric Cognitive Disorders*, 22(1), 108-20.

[Janata P. 2009. *The neural architecture of music-evoked autobiographical memories. Cereb Cortex*. 19\(11\),2579-94.](#)

Janata, P., Tomic, S.T., & Rakowski, S.K. 2007. Characterization of music-evoked autobiographical memories. *Memory*, 15, 845-860.

Katagiri, J. 2009. The effect of background music and song texts on the emotional understanding of children with autism. *Journal of Music Therapy*, 46, 15-31.

Kilgour, A.R., Jakobson, L.S., & Cuddy, L.L. 2000. Music training and rate of presentation as mediators of text and song recall. *Memory Cognition*, 28, 700-710.

Knox, R. & Jutai, J. 1996. Music-based rehabilitation of attention following brain injury. *Canadian Journal of Rehabilitation*, 9, 169-181.

Knox, R., Yokota-Adachi, H., Kershner, J., & Jutai, J. 2003. Musical attention training program and alternating attention in brain injury: An initial report. *Music Therapy Perspectives*, 21, 99-104.

Kostka, M. 1993. A comparison of selected behaviors of a student with autism in special education and regular music class. *Music Therapy Perspectives*, 11, 57-60.

Loui, P., Grent-'t-Jong, T., Torpey, D., Woldorff, M. 2005. Effects of attention on the neural processing of harmonic syntax in Western Music. *Cognitive Brain Research*, 25 (3), 678-87.

Ma, Y., Nagler, J., Lee, M., & Cabrera, I. 2001. Impact of music therapy on the communication skills of toddlers with pervasive developmental disorder. *Annals of the New York Academy of Sciences*, 930, 445-447.

Maeller, D.H. 1996. Rehearsal strategies and verbal working memory in multiple sclerosis. Unpublished master thesis, Colorado State University.

Magee, W.L. 2007. Music as a diagnostic tool in low awareness states: Considering limbic responses. *Brain Injury*, 21, 593-599.

Magee W.L. 2005. Music therapy with patients in low awareness states: approaches to assessment and treatment in multidisciplinary care. *Neuropsychol Rehabil.* 15(3-4), 522-36.

Mammarella, N., Fairfield, B., Cornoldi C. 2007. Does music enhance cognitive performance in healthy older adults? The Vivaldi Effect. *Ageing Clinical and Experimental Research*, October. [E-Publication]

Manly, T., Hawkins, K., Evans, J., Woldt, K., Robertson, I.H. 2002. Rehabilitation of executive function: facilitation of effective goal management on complex tasks using periodic auditory alerts. *Neuropsychologia*, 40(3), 271-81.

Marques, C., Moreno, S., Luis Castro, S., Besson, M. 2007. *Journal of Cognitive Neuroscience*, 19(9), 1453-63.

McCombe, W.S., Harris-Love, M., Liu, W., & Whittall, J. 2006. Temporal coordination of the arms during bilateral simultaneous and sequential movements in patients with chronic hemiparesis. *Experimental Brain Research*, 168(3), 450-4.

McGivern, R.F., Berka, C., Languis, M.L., & Chapman, D. 1991. Detection of deficits in temporal pattern discrimination using the Seashore rhythm test in young children with reading impairments. *Journal of Learning Disabilities*, 24, 58-62.

McIntosh, G.C., Peterson, D.A., & Thaut, M.H. 2006a. Verbal learning with a musical template increases neuronal synchronization and improves verbal memory in patients with Multiple Sclerosis (MS). *Proceedings of the 4th World Congress for Neurorehabilitation*, P1-018.

McIntosh, G.C., Peterson, D.A., & Thaut, M.H. 2006b. Verbal learning with a musical template increases neuronal synchronization and improves verbal memory in patients with multiple sclerosis. *Neurorehabilitation & Neural Repair*, 20, 129.

[Meilan JJ, Iodice R, Carro J, Sánchez JA, Palmero F, Mateos AM. 2011. Improvement of the autobiographic memory recovery by means of sad music in AD Type dementia. *Aging Clin Exp Res*. 2011 Jul 21. \[Epub ahead of print\]](#)

Millard, K.A.O. & Smith, J.A. 1989. The influence of group singing therapy on the behavior of Alzheimer's disease patients. *Journal of Music Therapy*, 26, 58-70.

Mitani, C., Nakata, T., Trehub, S.E., Kanda, Y., Kumagami, H., Takasaki, K., Miyamoto, I., Takahashi, H. 2007. *Ear and Hearing*, 28(2), 29S-33S.

Moore, K.S., Peterson, D.A., O'Shea, G., McIntosh, G.C., & Thaut, M.H. 2008. The effectiveness of music as a mnemonic device on recognition memory for people with multiple sclerosis. *Journal of Music Therapy*, 45, 307-329.

Morton, L.L., Kershner, J.R., & Siegel, L.S. 1990. The potential for therapeutic applications of music on problems related to memory and attention. *Journal of Music Therapy*, 26, 58-70.

Mottron, L., Peretz, I., & Menard, E. 2000. Local and global processing of music in high-functioning persons with autism: Beyond central coherence. *Journal of Child Psychology and Psychiatry*, 41 (8), 1057-1065.

Moussard, A. et al. (2012). Music as an aid to learn new verbal information in Alzheimer's disease. *Music Perception*, 29(5), 521-531.

Narme, P., Tonini, A., Khatir, F., Schiaratura, L., Clément, S., Samson, S., (2012). Non-pharmacological treatment for Alzheimer's disease: comparison between musical and non-

musical interventions. *Gériatrie et Psychologie Neuropsychiatrie du Vieillessement*. 10 (2), pp.215-224.

Nieuwboer, A., Kwakkel, G., Rochester, L., Jones, D., van Wegen, E., Willems, A.M., Chavret, F., Hetherington, V., Baker, K., & Lim, I. 2007. Cueing training in the home improves gait-related mobility in Parkinson's disease: the RESCUE trial. *Journal of Neurology, Neurosurgery & Psychiatry*, 78(2), 134-40.

Noda, R., Maeda, Y., & Yoshino, A. 2004. Therapeutic time window for musicokinetic therapy in a persistent vegetative state after severe brain damage. *Brain Injury*, 18 (5), 509-515.

Nayak, S. et al. (2000). Effect of music therapy on mood and social interaction among individuals with acute traumatic brain injury and stroke. *Rehabilitation Psychology*, 45(3), 274-283.

O'Callaghan, C.C. 1996. Pain, music creativity and music therapy in palliative care. *American Journal of Hospice & Palliative Care*, 13, 43-49.

O'Callaghan, C.C. 1993. Communicating with brain-impaired palliative care patients through music therapy. *Journal of Palliative Care*, 9, 53-55.

O'Shea, G., Sena, K.M., Peterson, D.A., Clegg, B.A., McIntosh, G.C., & Thaut, M.H. 2005. Representation without retention: Implicit learning in adults with multiple sclerosis. *Journal of Cognitive Neuroscience (suppl)*, Cognitive Neuroscience Society 12th Annual Meeting, In Press

Ozdemir, L. & Akdemir, N. 2009. Effects of multisensory stimulation on cognition, depression and anxiety levels of mildly-affected Alzheimer's patients. *Journal of the Neurological Sciences*. 283, 211-213.

Parbery-Clark, A., Skoe, E., Lam, C., & Kraus, N. 2009. Musician enhancement for speech-in-noise. *Ear Hear*, 30, 653-61.

[Peng SM](#), [Koo M](#), [Kuo JC](#). 2010. [Effect of group music activity as an adjunctive therapy on psychotic symptoms in patients with acute schizophrenia](#). *Arch Psychiatr Nurs*. 24(6),429-34.

Petersen B, Mortensen MV, Gjedde A, Vuust P. 2009. Reestablishing speech understanding through musical ear training after cochlear implantation: a study of the potential cortical plasticity in the brain. *Ann N Y Acad Sci*. 1169,437-40.

Peterson, D.A. & Thaut, M.H. 2007. Music increases frontal EEG coherence during verbal learning. *Neuroscience Letters*, 412, 217-221.

Purdie, H. 1997. Music therapy in neurorehabilitation: Recent developments and new challenges. *Critical Reviews in Physical and Rehabilitation Medicine*, 9, 205-217.

Quintin, E.M., Bhatara, A., Poissant, H., Fombonne, E., Levitin, D.J., (2010). Emotion perception in music in high-functioning adolescents with Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*, Dec. 22, 2010.

- Raglio, A. et al. (2012). Neuroscientific bases and effectiveness of music and music therapy in neuromotor rehabilitation. *Neurorehabilitation and Neural Repair*, 12(1611).
- Rainey, D. & Larsen, J. 2002. The effect of familiar melodies on initial learning and long-term memory for unconnected text. *Music Perception*, 20 (2), 173-186.
- Robb, S.L. (Ed.) 2003. *Music Therapy in Pediatric Healthcare: Research and Evidence-Based Practice*. Silver Spring, MD: American Music Therapy Association, Inc.
- Robertson, I.H., Manly T., Beschin, N., Daini, R., Haeske-Dewick, H., Homberg, V., Jehkonen, M., Pizzamiglio, G., Shiel, A., Weber, E. 1997. Auditory sustained attention is a marker of unilateral spatial neglect. *Neuropsychologia*, 35(12), 1527-32.
- Robin, D.A., Tranel, D. & Damasio, H. 1990. Auditory perception of temporal and spectral events in patients with focal left and right cerebral lesions. *Brain and Language*, 39, 539-555.
- Rosenbaum, R.S., Furey, M.L., Horwitz, B., & Grady, C.L. 2004. Altered communication between emotion-related brain regions supports short-term memory in Alzheimer' disease. *Proceedings Society for Neuroscience*, 203.8. [Abstract]
- Roy, M., Peretz, I., Rainville, P. 2007. Emotional valence contributes to music-induced analgesia. *Pain*, May. [E-Publication]
- Sacks, O. 1998. Music and the Brain. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 1-18). St. Louis, MO: MMB Music, Inc.
- Samson, S., Dellacherie, D. & Platel, H. 2009. Emotional power of music in patients with memory disorders. *Annals of New York Academy of Sciences*, 1169, 245-255.
- Särkämö, T. et al. (2010). Music and speech listening enhance the recovery of early sensory processing after stroke. *Journal of Cognitive Neuroscience*, 22(12), 2716-2727.
- Sarkomo, T., Tervaniemi, M., Latinen, S., Forsblom, A., Soynila, S., Mikkonen, M., Antti, T., Silvennoinen, H.M., Erkkila, J., Laine, M., Peretz, I., Hietanen, M. 2008. Music listening enhances cognitive recovery and mood after middle cerebral artery stroke. *Brain*. [E-Publication]
- Schweizer, S., Hampshire, A., Dalgleish, T., (2011). Extending Brain-Training to the Affective Domain: Increasing Cognitive and Affective Executive Control through Emotional Working Memory Training. *PLoS One*. 6 (9), E-page: 24372.
- Sears, W. 1968. Processes in music therapy. In E.T. Gaston (Ed.), *Music in Therapy* (pp. 30-44). New York, NY: MacMillan Co.
- Silverman, MJ. 2007. The effect of paired pitch, rhythm, and speech on working memory as measured by sequential digit recall. *Journal of Music Therapy*, Winter.
- Simmons-Stern NR, Deason RG, Brandler BJ, Frustace BS, O'Connor MK, Ally BA, Budson AE. 2012. Music-based memory enhancement in Alzheimer's Disease: Promise and limitations. *Neuropsychologia*. 2012 Sep 19. [Epub ahead of print]

- Son, G.R., Therrien, B., & Whall, A. 2002. Implicit memory and familiarity among elders with dementia. *Journal of Nursing Scholarship*, 34, 263-267.
- Soto, D., Funes, M.J., Guzman-Garcia, A., Warbick, T., Rotshtein, P., Humphreys, G.W., (2009). Pleasant music overcomes the loss of awareness in patients with visual neglect. *Proceedings of the National Academy of Sciences*. 106 (14), pp.6011-6016.
- Sussman, J.E. 2009. The effect of music on peer awareness in preschool age children with developmental disabilities. *Journal of Music Therapy*, 46, 53-68.
- Suzuki, M., Kanamori, M., Watanabe, M., Nagasawa, S., Kojima, E., Ooshiro, H., Nakahara, D. 2004. Behavioral and endocrinological evaluation of music therapy for elderly patients with dementia. *Nursing & Health Sciences*, 6(1), 11-8.
- Thaut, M.H. 1987. Visual versus auditory (musical) stimulus preferences in autistic children: A pilot study. *Journal of Autism and Developmental Disorders*, 17, 425-432.
- Thaut, M.H. 1988. Measuring musical responsiveness in autistic children: A comparative analysis of improvised musical tone sequences of autistic, normal, and mentally retarded individuals. *Journal of Autism and Developmental Disorders*, 18, 561-571.
- Thaut, M.H. 1989. Music therapy, affect modification, and therapeutic change: Towards an integrative model. *Music Therapy Perspectives*, 7, 55-62.
- Thaut, M.H. 1997. Wer und was fordert die Musiktherapie in der Zukunft? Wissenschaftstheoretische Grundlagen, Forschungsmodelle un therapeutische Anwendungen. *Musiktherapeutische Umschau*, 18, 15-24. [In German]
- Thaut, M.H. 1998. Fondamenti Scientifici Della Musica. In *Terapia: Una Nuova Cornice Teorica*. In G.G. Perilli & F. Russo (Eds.), *La Medicina Dei Suoni* (pp. 94-111). Rome, Italy: Edizioni Borla. [In Italian]
- Thaut, M.H. 1999. Einfuehrung in die neurologische Musiktherapie. *Neurologie und Rehabilitation*, 5 (suppl 2), 11. [In German] [Abstract]
- Thaut, M.H. & Peterson, D.A. 2000. Event related potential correlates of rhythmic auditory synchronization. *Proceedings Society for Neuroscience*, 833.6. [Abstract]
- Thaut, M.H. & Mahraun. 2004. The influence of music and rhythm on a sustained attention task in children with autism. Submitted for publication.
- Thaut, M.H., Nickel, A., & Hoemberg, V. 2004. Neurologic music therapy: Compendium of scientific background and clinical methodology. *Musiktherapeutische Umschau*, 25, 35-44. [In German]
- Thaut, M.H., Peterson, D.A., & McIntosh, G.C. 2005. Temporal entrainment of cognitive function: Musical mnemonics induce brain plasticity and oscillatory synchrony in neural networks underlying memory. *Annals of the New York Academy of Sciences*, 1060, 243-254.
- Thaut, M.H., Gardiner, J.C., Holmberg, D., Horwitz, J., Kent, L., Andrews, G., Donelan, B., McIntosh, G.R. 2009. Neurologic music therapy improves executive function and emotional

adjustment in traumatic brain injury rehabilitation. *Neurosciences and music III- Disorders and plasticity*, Ann. N.Y. Acad. Sci., xxxx, 1-11.

Thaut, M.H., Sena, K.M., McIntosh, G.C., & Peterson, D.A. 2007. Musical structure facilitates verbal learning in multiple sclerosis. *Musical Structure Facilitates*, 325-330.

Tomaino, C.M. 1998. Music and memory: accessing residual function. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 19-27). St. Louis, MO: MMB Music, Inc.

[Ulrich G, Houtmans T, Gold C.](#) 2007. [The additional therapeutic effect of group music therapy for schizophrenic patients: a randomized study.](#) *Acta Psychiatr Scand.* 116(5),362-70.

Unkefer, R.F. & Thaut, M.H. (Eds.). 2002. *Music Therapy in the Treatment of Adults with Mental Disorders* (2nd ed.). St. Louis, MO: MMB Music, Inc.

Van Besouw, R.M., Howard, D.M., Ternstrom, S. 2005. Towards an understanding of speech and song perception. *Logopedics, Phoniatrics, Vocology*, 30(3-4), 129-35.

Van Wegen, E., de Goede, C., Lim, I., Rietberg, M., Nieuwboer, A., Willems, A., Jones, D., Rochester, L., Hetherington, V., Berendse, H., Zijlmans, J., Wolters, E., & Kwakkel, G. 2006. The effect of rhythmic somatosensory cueing on gait in patients with Parkinson's disease. *Journal of the Neurological Sciences*, 248(1-2), 210-4.

Van Wegen, E., Lim, I., de Goede, C., Nieuwboer, A., Willems, A., Jones, D., Rochester, L., Hetherington, V., Berendse, H., Zijlmans, J., Wolters, E., & Kwakkel, G. 2006. The effects of visual rhythms and optic flow on stride patterns of patients with Parkinson's disease. *Parkinsonism & Related Disorders*, 12(1), 21-7.

Vanstone, A.D. & Cuddy, L.L. 2010. Musical Memory in Alzheimer disease. *Aging, Neuropsychology, and Cognition*, 17, 108-128.

Vanstone, A.D., Cuddy, L.L., Duffin, J.M., & Alexander, E. 2009. Exceptional preservation of memory for tunes and lyrics: Case studies of amusia, profound deafness, and Alzheimer's disease. *Annals of the New York Academy of Sciences*, 1169, 291-294.

Wallace, W.T. 1994. Memory for music: Effect of melody on recall of text. *Journal of Experimental Psychology: Learning, Memory, Cognition*, 20, 1471-1485.

Wheeler, B. et al. (2003). Effects of number of sessions and group or individual music therapy on the mood and behavior of people who have had strokes or traumatic brain injuries. *Nordic Journal of Music Therapy*, 12(2), 139-151.

Willems, A.M., Nieuwboer, A., Chavret, F., Desloovere, K., Dom, R., Rochester, L., Jones, D., Kwakkel, G., & Van Wegen, E. 2006. The use of rhythmic auditory cues to influence gait in patients with Parkinson's disease, the differential effect for freezers and non-freezers, an explorative study. *Disability & Rehabilitation*, 28(11), 721-8.

Wimpory, D., Chadwick, P., & Nash, S. 1995. Brief report: Musical interaction therapy for children with autism: An evaluative case study with two-year follow-up. *Journal of Autism and Developmental Disabilities*, 25 (3), 231-248.

Wit, V., Knox, R., Jutai, J., & Loveszy, R. 1994. Music therapy and rehabilitation of brain injury: A pilot study. *Canadian Journal of Music Therapy*, 2, 72-89.

Wolfe, D.E. & Hom, C. 1993. Use of melodies as structural prompts for learning and retention of sequential verbal information by preschool students. *Journal of Music Therapy*, 30, 100-118.

Wolfe, D.E. & Noguchi, L.K. 2009. The use of music with young children to improve sustained attention during a vigilance task in the presence of auditory distractions. *Journal of Music Therapy*, 46, 69-83.

Yannou, D., Baloyiannis, S., Kostarido- Eukleides, A., Kioseoglou, G., Damianou- Marinis, E. Klonari, D., nakopoulou, E., Tzedaki, K. 2003. Music Perception in Patients with Dementia due to Alzheimer's Disease. *Proceedings of the fifth triennial ESCOM conference*, 548-551.