

SENSORIMOTOR

Clinical Research

Abiru, M., Kikuchi, Y., Tokita, K., Mihara, Y., Fujimoto, M., Mihara, B. 2007. Neurologic Music Therapy for Rehabilitation for Stroke Patients-Prospects of Music Therapy for Gait Disturbance. *Neurological Therapeutics*, 24(6), 711-718.

Altenmüller E, Marco-Pallares J, Münte TF, Schneider S. (2009). Neural reorganization underlies improvement in stroke-induced motor dysfunction by music-supported therapy. *Ann N Y Acad Sci*. Jul(1169),395-405.

Altenmuller, E., Marco-Pallares, J., Munte, T.F. & Schneider, S. 2009. Neural reorganization underlies improvement in stroke-induced motor dysfunction by music-supported therapy. *Annals of the New York Academy of Sciences*, 1169, 395-405.

Arias, P., Cudeiro, J. (2008) Effects of rhythmic sensory stimulation (auditory, visual) on gait in Parkinson's disease patients. *Exp Brain Res*. 186 (4), 589-601.

Baram Y, Lenger R. 2012. Gait improvement in patients with cerebral palsy by visual and auditory feedback. *Neuromodulation*. 15(1),48-52.

Baram, Y. & Miller, A. 2007. Auditory feedback control for improvement of gait in patients with multiple sclerosis. *Neurological Sciences*, 254, 90-94.

Bernardi NF, Aggujaro S, Caimmi M, Molteni F, Maravita A, Luzzatti C. 2009. A new approach to rhythm cueing of cognitive functions: the case of ideomotor apraxia. *Ann N Y Acad Sci*. July(1169),417-21.

Bernatzky, G., Bernatzky, P., Hesse, H.P., Staffen, W., & Ladurner, G. 2004. Stimulating music increases motor coordination in patients afflicted by Morbus Parkinson. *Neuroscience Letters*, 361, 4-8.

Bryant MS, Rintala DH, Lai EC, Protas EJ. 2009. An evaluation of self-administration of auditory cueing to improve gait in people with Parkinson's disease. *Clinical Rehabilitation*. 23(12), 1078-1085.

Buetefish, C., Hummelsheim, H., Denzler, P., & Mauritz, K.H. 1995. Repetitive training of isolated movements improves the outcome of motor rehabilitation of the centrally paretic hand. *Journal of Neurological Sciences*, 130, 59-68.

Carver, F. W., Fuchs, A., Jantzen, K. J., & Kelso, J. (2002). Spatiotemporal analysis of the neuromagnetic response to rhythmic auditory stimulation: Rate dependence and transient to steady-state transition. *Clinical Neurophysiology*,113(12), 1921-1931

Clair AA, O'Konski M. (2006). The effect of rhythmic auditory stimulation (RAS) on gait characteristics of cadence, velocity, and stride length in persons with late stage dementia. *J Music Ther*. 43(2), 154-63.

Clark, C. & Chadwick, D. 1980. Clinically Adapted Instruments for the Multiply Handicapped. St. Louis, MO: Magnamusic-Baton.

Conklyn, D., Stough, D., Novak, E., Paczak, S., Chemali, K., & Bethoux, F. (2010). A home-based walking program using rhythmic auditory stimulation improves gait performance in patients with multiple sclerosis: A pilot study. *Neurorehabilitation And Neural Repair*, 24(9), 835-842

Cross, P., McLellan, M., Vomberg, E., Monga, M., & Monga, T.N. 1984. Observations on the use of music in rehabilitation of stroke patients. *Physiotherapy Canada*, 36, 197-201.

[Cubo E](#), [Leurgans S](#), [Goetz CG](#). 2004. Short-term and practice effects of metronome pacing in Parkinson's disease patients with gait freezing while in the 'on' state: randomized single blind evaluation. *Parkinsonism & Related Disorders*. 10(8); 507-510.

Davis, W.B., Gfeller, K., & Thaut, M.H. 1992 (1998, 2nd ed.). An Introduction to Music Therapy: Theory and Practice. Dubuque, IA: Wm. C. Brown Publishers. (Japanese Language Edition, 1997. Sapporo, Japan: Ichibaku Shuppansya Publishing.) (Spanish Language Edition, 2000) (Korean Language Edition, 2002)

[de Bruin N](#), [Doan JB](#), [Turnbull G](#), [Suchowersky O](#), [Bonfield S](#), [Hu B](#), [Brown LA](#). 2010. Walking with music is a safe and viable tool for gait training in Parkinson's disease: the effect of a 13-week feasibility study on single and dual task walking. *Parkinsons Disorders*. Jul 13 2010, 483530.

[de Dreu MJ](#), [van der Wilk AS](#), [Poppe E](#), [Kwakkel G](#), [van Wegen EE](#). 2012. Rehabilitation, exercise therapy and music in patients with Parkinson's disease: a meta-analysis of the effects of music-based movement therapy on walking ability, balance and quality of life. *Parkinsonism & Related Disorders*. 18(Suppl 1), S114-119.

de l'Etoile SK. 2008. The effect of rhythmic auditory stimulation on the gait parameters of patients with incomplete spinal cord injury: an exploratory pilot study. *Int J Rehabil Res* 31(2), 155-7.

Del Olmo, Aria P., Furio, M., Pozo, M., & Cudeiro, J. 2006. Evaluation of the effect of training using auditory stimulation on rhythmic movement in Parkinsonian patients—a combined motor and [18F]-FDG PET study. *Parkinsonism & Related Disorders*, 12(3), 155-64.

Del Olmo, M.F., & Cudeiro, J. 2003. A simple procedure using auditory stimuli to improve movement in Parkinson's disease: A pilot study. *Neurology & Clinical Neurophysiology*, 2003 (2), 1-7.

Del Olmo, M.F., & Cudeiro, J. 2003. The timing in Parkinson's disease: Effects of a rehabilitation program based on rhythmic sound cues. *Proceedings Society for Neuroscience*, 734.2.

Dozza, M., Chiari, L., Hlavacka, F., Cappello, A., & Horak, F.B. 2006. Effects of linear versus sigmoid coding of visual or audio biofeedback for the control of upright stance. *IEEE Transactions on Neural Systems & Rehabilitation Engineering*, 14(4), 505-12.

Dozza, M., Chiari, L., & Horak, F.B. 2005. Audio-biofeedback improves balance in patients with bilateral vestibular loss. *Archives of Physical Medicine and Rehabilitation*, 86 (7), 1401-3.

Elliott, B. 1982. Guide to the Selection of Musical Instruments with Respect to Physical Ability and Disability. Saint Louis, MO: MMB Music, Inc.

Fields, B. 1954. Music as an adjunct in the treatment of brain damaged patients. *American Journal of Physical Medicine*, 33, 273-283.

Ford, M., Wagenaar, R., Newell, K. 2007. The effects of auditory rhythms and instruction on walking patterns in individuals post stroke. *Gait and Posture*, 26, 150-155.

Frazzitta, G., Maestri, R., Uccellini, D., Bertotti, G., Abelli, P. 2009. Rehabilitation treatment of gait in patients with Parkinson's disease with freezing: a comparison between two physical therapy protocols using visual and auditory cues with or without treadmill training. *Movement Disorders*. 24(8), 1139-43.

Freedland, R.L., Festa, C., Sealy, M., McBean, A., Elghazaly, P., Capan, A., Brozycki, L., Nelson, A.J., & Rothman, J. 2002. The effects of pulsed auditory stimulation on various gait measurements in persons with Parkinson's disease. *Neurorehabilitation*, 17 (1), 81-87.

Freeman, J.S., Cody, F.W., & Schady, W. 1993. The influence of external timing cues upon the rhythm of voluntary movements in Parkinson's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*, 56, 1078-1084.

Fujioka T, Ween JE, Jamali S, Stuss DT, Ross B. 2012. Changes in neuromagnetic beta-band oscillation after music-supported stroke rehabilitation. *Ann N Y Acad Sci*. Apr (1252), 294-304.

Gaston, E.T. (Ed.). 1968. Music in Therapy. New York, NY: MacMillan Co.

Gfeller, K.E. 1999. Music therapy in the treatment of sensory disorders. In W.B. Davis, K.E. Gfeller, & M.H. Thaut (Eds.), *An Introduction to Music Therapy*. Boston, MA: McGraw Hill.

Goldshtröm Y, Knorr G, Goldshtröm I. 2010. Rhythmic exercises in rehabilitation of TBI patients: a case report. *J Bodyw Mov Ther*. 14(4), 336-45.

Hausdorff J.M., Lowenthal, J., Herman, T., Gruendlinger, L., Peretz, C., & Giladi, N. (2007) Rhythmic auditory stimulation modulates gait variability in Parkinson's disease. *European Journal of Neuroscience*. 26 (8), 2369-2375.

Hesse, S. & Werner, C., & Bardeleben, A. 2004. Electromechanical gait training with functional electrical stimulation: Case studies in spinal cord injury. *Spinal Cord*, 42, 346-52.

Hesse, S. & Werner, C. 2003. Post-stroke motor dysfunction and spasticity: Novel pharmacological and physical treatment strategies. *CNS Drugs*, 17, 1070-1093.

Hove, M. et al. (2012). Interactive rhythmic auditory stimulation reinstates natural 1/f timing in gait of Parkinson's patients. *PLoS ONE*, 7(3).

Howe, T.E., Lovgreen, B., Cody, F.W., Ashton, V.J., & Oldham, J.A. 2003. Auditory cues can modify the gait of persons with early-stage Parkinson's disease: A method for enhancing Parkinsonian walking performance? *Clinical Rehabilitation*, 17 (4), 363-367.

Hurt-Thaut, C.P. & Johnson, S.B. 2003. Neurologic music therapy with children: Scientific foundations and clinical applications. In S.L. Robb (Ed.), *Music Therapy in Pediatric Healthcare: Research and Evidence-Based Practice* (pp. 81-100). Silver Spring, MD: American Music Therapy Association, Inc.

Hurt, C.P., Rice, R.R., McIntosh, G.C., & Thaut, M.H. 1998. Rhythmic auditory stimulation in gait training for patients with traumatic brain injury. *Journal of Music Therapy*, 35, 228-241.

Jeong, S., Kim, M.T. 2007. Effects of a theory-driven music and movement program for stroke survivors in a community setting. *Applied Nursing Research*, 20(3), 125-31.

Johannsen L, Wing AM, Pelton T, Kitaka K, Zietz D, Brittle N, van Vliet P, Riddoch J, Sackley C, McManus R. 2010. Seated bilateral leg exercise effects on hemiparetic lower extremity function in chronic stroke. *Neurorehabilitation and Neural Repair*. 24(3), 243-53.

Kadivar, Z., Corcos, D. M., Foto, J., & Hondzinski, J. M. (2011). Effect of step training and rhythmic auditory stimulation on functional performance in Parkinson patients. *Neurorehabilitation And Neural Repair*, 25(7), 626-635.

Kaipust JP, McGrath D, Mukherjee M, Stergiou N. 2012. Gait Variability is Altered in Older Adults When Listening to Auditory Stimuli with Differing Temporal Structures. *Ann Biomed Eng*. Sep 7. [Epub ahead of print]

Kenyon, G.P. & Thaut, M.H. 2000. A measure of kinematic limb instability modulation by rhythmic auditory stimulation. *Journal of Biomechanics*, 33, 1313-1323.

Kim, S.J., Kwak, E.E., Park, E.S., Lee, D.S., Kim, K.J., Song, J.E., Cho, S.R., (2011). Changes in gait patterns with rhythmic auditory stimulation in adults with cerebral palsy. *NeuroRehabilitation*. 29 (3), pp.233-241.

Kim JS, Oh DW, Kim SY, Choi JD. 2011. Visual and kinesthetic locomotor imagery training integrated with auditory step rhythm for walking performance of patients with chronic stroke. *Clin Rehabil*. 25(2),134-45.

Kimber, T.E., Tsai, C.S., Semmler, J., Brophy, B.P., & Thompson, P.D. 1999. Voluntary movement after pallidotomy in severe Parkinson's disease. *Brain*, 122, 895-906.

Krasovsky T, Berman S, Liebermann DG. 2010. Kinematic features of continuous hand reaching movements under simple and complex rhythmical constraints. *Journal of Electromyography and Kinesiology*. 20(4), 636-41.

Kwak, E.E. 2007. Effect of rhythmic auditory stimulation on gait performance in children with spastic cerebral palsy. *Journal of Music Therapy*, 44, 198-216.

Lamontagne, A. & Fung J. 2004. Faster is better: Implications for speed-intensive gait training after stroke. *Stroke*, 35 (11), 2543-8.

Ledger, S., Galvin, R., Lynch, D., Stokes, E.K. 2008. A randomised controlled trial evaluating the effect of an individual auditory cueing device on freezing and gait speed in people with Parkinson's disease. *BMC Neurology*. 8, 46.

L'Etoile, S.K. 2008. The effect of rhythmic auditory stimulation on the gait parameters of patients with incomplete spinal cord injury: An exploratory pilot study. *International Journal of Rehabilitation Research*, 31, 155-157.

Liberzon, T. & Brown, S.H. 1998. Effects of rhythmic auditory cueing on timing variability of sequential arm movements in Parkinson's disease. *Proceedings Society for Neuroscience*, 653.18.

Lim HA, Miller K, Fabian C. 2011. The effects of therapeutic instrumental music performance on endurance level, self-perceived fatigue level, and self-perceived exertion of inpatients in physical rehabilitation. *J Music Ther*. 48(2),124-48.

Lim, I., Van Wegen, E., de Goede, C., Deutekom, M., Nieuwboer, A., Willems, A., & Jones, D. 2005. Effects of external rhythmical cueing on gait in patients with Parkinson's disease. *Clinical Rehabilitation*, 19 (7), 695-713.

Lohnes, CA, Earhart, GM. 2011. The impact of attentional, auditory, and combined cues on walking during single and cognitive dual tasks in Parkinson disease. *Gait & Posture*. 33 (3), 478-83.

Luft, A.R., McCombe-Waller, S., Whitall, J., Forrester, L.W., Macko, R., Sorkin, J.D., Schulz, J.B., Goldberg, A.P., & Hanley, D.F. 2004. Repetitive bilateral arm training and motor cortex activation in chronic stroke: A randomized controlled trial. *JAMA*, 292 (15), 1853-1861.

Ma, H.I., Hwang, W.J., & Lin, K.C. 2009. The effects of two different auditory stimuli on functional arm movement in persons with Parkinson's disease: a dual-task paradigm. *Clinical Rehab*, 23, 229-237.

Mak, M. (2006). Feed-forward audio-visual cues could enhance sit-to-stand in Parkinsonian patients. *Proceedings of the 4th World Congress for Neurorehabilitation*, F1B-7.

Malcolm, M.P., Massie, C., Thaut, M.H. 2009. Rhythmic auditory- motor entrainment improves hemiparetic arm kinematics during reaching movements: A pilot study. *Top Stroke Rehabil*, 16, 69-79.

Mandel, A.R., Nymark, J.R., Balmer, S.J., Grinnell, D.M., & O'Riain, M.D. 1990. Electromyographic feedback versus rhythmic positional biofeedback in computerized gait retraining with stroke patients. *Archives of Physical and Medical Rehabilitation*, 71, 649-654.

Mauritz, K.H. 2002. Gait training in hemiplegia. *European Journal of Neurology*, 9 suppl 1:23-29; discussion 53-61.

McCombe, W.S. & Whitall, J. 2005. Hand dominance and side of stroke affect rehabilitation in chronic stroke. *Clinical Rehabilitation*, 19 (5), 544-551.

McCombe Waller, S. & Whitall, J. (2006). Combining bilateral and distal arm training to promote arm and hand recovery in patients with chronic hemiparesis: A case report. Proceedings of the 4th World Congress for Neurorehabilitation., P2-080

McIntosh, G.C., Brown, S.H., Rice, R.R., & Thaut, M.H. 1997. Rhythmic auditory-motor facilitation of gait patterns in patients with Parkinson's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*, 62, 122-126.

McIntosh, G.C., Rice, R.R., Hurt, C.P., & Thaut, M.H. 1998. Long-term training effects of rhythmic auditory stimulation on gait in patients with Parkinson's disease. *Movement Disorders*, 13 (2), 212. [Abstract]

McIntosh, G.C., Rice, R.R., Thaut, & M.H. 1994. Stride frequency modulation in Parkinsonian gait using rhythmic auditory stimulation. *Annals of Neurology*, 36, 316. [Abstract]

McIntosh, G.C. & Thaut, M.H. 1997. Rhythmic sensorimotor facilitation of gait in patients with Parkinson's disease. *Proceedings of the 4th Australian Multidisciplinary Conference on Parkinson's Disease*, 63-65. [Abstract]

McIntosh, G.C., Thaut, M.H., & Rice, R.R. 1996. Rhythmic auditory stimulation as entrainment and therapy technique in gait of stroke and Parkinson's disease patients. In R. Pratt & R. Spintge (Eds.), *Music Medicine*, Vol. II (pp. 145-152). St. Louis, MO: MMB Music.

McIntosh, G.C., Thaut, M.H., Rice, R.R., Miller, R.A., Rathbun, J., & Brault, J.M. 1995. Rhythmic facilitation in gait training of Parkinson's disease. *Annals of Neurology*, 38, 331. [Abstract]

McIntosh, G.C., Thaut, M.H., Rice, R.R., & Prassas, S.G. 1993. Auditory rhythmic cuing in gait rehabilitation with stroke patients. *Canadian Journal of Neurological Sciences*, 20, 168. [Abstract]

McIntosh, G.C., Thaut, M.H., Rice, R.R., & Prassas, S.G. 1995. Rhythmic facilitation of gait kinematics in stroke patients. *Journal of Neurologic Rehabilitation*, 9, 131. [Abstract]

Miller, R.A., Thaut, M.H., McIntosh, G.C., Rice, R.R. 1996. Components of EMG symmetry and variability in Parkinsonian and healthy elderly gait. *Electroencephalography and Clinical Neurophysiology*, 101, 1-7.

Mitoma H, Yoneyama M, Orimo S. 2010. 24-hour recording of parkinsonian gait using a portable gait rhythmogram. *Intern Med*. 49(22),2401-8.

Molinari, M., Leggio, M.G., Filippini, V., Cioia, M.C., Cerasa, A., & Thaut, M.H. 2005. Sensorimotor transduction of time information is preserved in subjects with cerebellar damage. *Brain Research Bulletin*, 67, 448-458.

Morris, G.S., Suteerawattananon, M., Etnyre, B.R., Jankovic, J., & Protas, E.J. 2004. Effects of visual and auditory cues on gait in individuals with Parkinson's disease. *Journal of the Neurological Sciences*, 219, 63-69.

Munneke, M., Keus, S.H.J., Bredero-Cohen, A.B., Hendriks, H.J., & Bloem, B.R. (2006). Evidence-based clinical practice guideline for physical therapy in Parkinson's disease. Proceedings of the 4th World Congress for Neurorehabilitation, P1-058.

[Nanhoe-Mahabier W](#), [Delval A](#), [Snijders AH](#), [Weerdesteyn V](#), [Overeem S](#), [Bloem BR](#). 2012. The possible price of auditory cueing: influence on obstacle avoidance in Parkinson's disease. *Movement Disorders*. 27(4), 574-578.

Nieuwboer, A., Kwakkel, G., Rochester, L., Jones, D., van Wegen E., Willems, A.M., Chavret, F., Hetherington, V., Baker, K., & Lim, I. 2007. Cueing training in the home improves gait-related mobility in Parkinson's disease: the RESCUE trial. *Journal of Neurology, Neurosurgery & Psychiatry*, 78(2), 134-40.

Pacchetti, C., Aglieri, R., Mancini, F., Martignoni, E., & Nappi, G. 1998. Active music therapy and Parkinson's disease: Methods. *Functional Neurology*, 13 (1), 57-67.

Pacchetti, C., Mancini, F., Aglieri, R., Fundaro, C., Martignoni, E., & Nappi, G. 2000. Active music therapy in Parkinson's disease: An integrative method for motor and emotional rehabilitation. *Psychosomatic Medicine*, 62 (3), 386-393.

Peng, Y.C., Lu, T.W., Wang, T.H., Chen, Y.L., Liao, H.F., Lin, K.H., Tang, P.F., (2011). Immediate effects of therapeutic music on loaded sit-to-stand movement in children with spastic diplegia. *Gait & Posture*. 33 (2), pp.274-278.

Picelli A, Camin M, Tinazzi M, Vangelista A, Cosentino A, Fiaschi A, Smania N. 2010. Three-dimensional motion analysis of the effects of auditory cueing on gait pattern in patients with Parkinson's disease: a preliminary investigation. *Neurol Sci*. 31(4),423-30.

Platz, T., Brown, R.G., & Marsden, C.D. 1998. Training improves the speed of aimed movements in Parkinson's disease. *Brain*, 121, 505-514.

Prassas, S.G. & Thaut, M.H. 1992. Effects of music and rhythm on gait characteristics of normal individuals. *Journal of Biomechanics*, 25, 684. [Abstract]

Prassas, S.G., Thaut, M.H., McIntosh, G.C., & Rice, R.R. 1997. Effect of auditory rhythmic cuing on gait kinematic parameters in stroke patients. *Gait and Posture*, 6, 218-223.

Rice, R.R., Thaut, M.H., McIntosh, G.C., & Miller, R.A. 1995. The effect of a home based gait training program for Parkinson's disease patients using rhythmic auditory stimulation. *Proceedings of the 12th International Congress of the World Federation for Physical Therapy*, 768. [Abstract]

Richards, C.L., Malouin, F., Bedard, P.J., & Cioni, M. 1992. Changes induced by L-Dopa and sensory cues on the gait of parkinsonian patients. In M. Woollacott and F. Horak (Eds.), *Posture and Control Mechanisms* (Vol. 2) (pp. 126-129). Eugene, OR: University of Oregon Books.

Robb, S.L. (Ed.) 2003. *Music Therapy in Pediatric Healthcare: Research and Evidence-Based Practice*. Silver Spring, MD: American Music Therapy Association, Inc.

- Robertson, S.D., Chua, R., Maraj, B.K., Kao, J.C., & Weeks, D.J. 2002a. Bimanual coordination dynamics in adults with down syndrome. *Motor Control*, 6, 388-407.
- Robertson, S.D., Van Gemmert, A.W., & Maraj, B.K. 2002b. Auditory information is beneficial for adults with down syndrome in a continuous bimanual task. *Acta Psychologica*, 110, 213-229.
- Rochester, L., Hetherington, V., Jones, D., Nieuwboer, A., Willems, A.M., Kwakkel, G., & Van Wegen, E. 2005. The effect of external rhythmic cues (auditory and visual) on walking during a functional task in homes of people with Parkinson's disease. *Archives of Physical Medicine & Rehabilitation*, 86 (5), 999-1006.
- Roerdink, M. et al. (2011). Walking to the beat of different drums: Practical implications for the use of acoustic rhythms in gait rehabilitation. *Gait and Posture*, 33(1), 690-694.
- Roerdink, M., Lamoth, C.J.C, van Kordelaar, J., Elich, P., Konijnenbelt, M., Kwakkel, G. & Beek, P.J. 2009. Rhythm perturbations in acoustically paced treadmill walking after stroke. *Neurorehabilitation and Neural Repair*, 23, 668-678.
- Roerdink, M., Lamoth, C.J.C, Kwakkel, G., van Wieringen, P.C.W. & Beek, P.J. 2007. Gait coordination after stroke: Benefits of acoustically paced treadmill walking. *Physical Therapy*, 87, 1009-1022.
- Sandrinini, G., Tassorelli, C., Balloni, L., Buscone, S., & Pacchetti, C. (2006). Efficacy of sensorial cues on gait rehabilitation in Parkinson's disease. *Proceedings of the 4th World Congress for Neurorehabilitation*, P1-053.
- Satoh, M. & Kuzuhara, S. 2008. Training in mental singing while walking improves gait disturbance in Parkinson's disease patients. *European Neurology*, 60, 237-243.
- Secoli, R. et al. (2011). Effect of visual distraction and auditory feedback on patient effort during robot-assisted movement training after stroke. *Journal of Neuroengineering and Rehabilitation*, 8(21).
- Senesac, Davis, and Richards. (2010). Generalization of a modified form of repetitive rhythmic bilateral training in stroke. *Human Movement Science*, 26(1), 137-148.
- Schauer, M. & Mauritz, K.H. 2003. Musical motor feedback (MMF) in walking hemiparetic stroke patients: Randomized trials of gait improvement. *Clinical Rehabilitation*, 17, 713-722.
- Schauer, M.L., Steingrueber, W., & Mauritz, K.H. 1996. Die Wirkung von Musik auf die Symmetrie des Gehens von Schlaganfallpatienten auf dem Laufband. *Biomedizinische Technik*, 41, 291-296.
- Schneider, S. et al. (2010). Music-supported training is more efficient than functional motor training for recovery of fine motor skills in stroke patients. *Music Perception*, 27(4), 271-280.
- Sears, W. 1968. Processes in music therapy. In E.T. Gaston (Ed.), *Music in Therapy* (pp. 30-44). New York, NY: MacMillan Co.

Schneider S, Schoenle PW, Altenmueller E, Munte TF. 2007. Using musical instruments to improve motor skill recovery following a stroke. *Journal of Neurology*, 254, 1339-1346.

Styns, F., van Noorden, L., Moelants, D., Moelants, D., Leman, M. 2007. Walking on music. *Human Movement Science*, September. [E-Publication]

Szmedra, L. & Bacharach, D.W. 1998. Effect of music on perceived exertion, plasma lactate, norepinephrine and cardiovascular hemodynamics during treadmill running. *International Journal of Sports Medicine*, 19, 32-37.

Teasell, R.W., Bhogal, S.K., Foley, N.C., & Speechley, M.R. 2003. Gait retraining post stroke. *Topics in Stroke Rehabilitation*, 10 (2), 34-65.

Thaut, M.H. 1985. The use of auditory rhythm and rhythmic speech to aid temporal muscular control in children with gross motor dysfunction. *Journal of Music Therapy*, 22, 108-128.

Thaut, M.H. 1988. Rhythmic intervention techniques in music therapy with gross motor dysfunction. *Arts in Psychotherapy*, 15, 127-137.

Thaut, M.H. 1992. Music in motor recovery with neurological disorders. Hearing before the Special Committee on Aging, United States Senate, 102nd Congress, August 1991 (pp. 100-101). Washington DC: U.S. Government Printing Office.

Thaut, M.H. 1996. Music therapy in neurologic rehabilitation. In H. Smeijsters & F. Mecklenbeck (Eds.), *Book of Abstracts, The 8th World Congress of Music Therapy* (p. 101). Bremen, Germany: Trialog Verlag.

Thaut, M.H. 1997. Clinical effects and neurological mechanisms of musical stimuli in neurological rehabilitation of elderly patients. *Proceedings of the Institute on Music Therapy and Aging*. Silver Spring, MD: National Association for Music Therapy, 13-14. [Abstract]

Thaut, M.H. 1997. Rhythmic auditory stimulation in rehabilitation of movement disorders: A review of current research. In D.J. Schneck & J.K. Schneck (Eds.), *Music in Human Adaptation* (pp. 223-230). Blacksburg, VA: Virginia Polytechnic Institute and State University.

Thaut, M.H., Hoemberg, V., Kenyon, G., & Hurt, C.P. 1998. Rhythmic entrainment of hemiparetic arm movements in stroke patients. *Proceedings Society for Neuroscience*, 653.7. [Abstract]

Thaut, M.H., Hurt, C.P., Dragan, D., & McIntosh, G.C. 1998. Rhythmic entrainment of gait patterns in children with cerebral palsy. *Developmental Medicine and Child Neurology*, 40 (78), 15. [Abstract]

Thaut, M.H., Hurt, C.P., & McIntosh, G.C. 1997. Rhythmic entrainment of gait patterns in traumatic brain injury rehabilitation. *Journal of Neurologic Rehabilitation*, 11, 131. [Abstract]

Thaut, M.H., Kenyon, G.P., Hurt, C.P., & Hoemberg, V. 1998. Rhythmic entrainment of

spatiotemporal parameters in paretic arm reaching movements of stroke patients. *Annals of Neurology* (submitted). [Abstract]

Thaut, M.H., Kenyon, G.P., Hurt, C.P., McIntosh, G.C., & Hoemberg, V. 2002. Kinematic optimization of spatiotemporal patterns in paretic arm training with stroke patients. *Neuropsychologia*, 40, 1073-1081.

Thaut, M.H., Kenyon, G.P., Schauer, M.L., & McIntosh, G.C. 1999. The connection between rhythmicity and brain function. *IEEE Engineering in Medicine and Biology Magazine*, 18 (2), 101-108.

Thaut, M.H., Lange, H., Miltner, R., Hurt, C.P., & Hoemberg, V. 1996. Rhythmic entrainment of gait patterns in Huntington's disease patients. *Proceedings Society for Neuroscience*, 727.6. [Abstract]

Thaut, M.H., Lange, H., Miltner, R., Hurt, C.P., & Hoemberg, V. 1998. Gait pattern analysis during velocity modulation with and without rhythmic sensory cues in patients with Huntington's disease. *Movements Disorders*, 13 (suppl 2), 157. [Abstract]

Thaut, M.H. & McIntosh, G.C. 1992. Effect of auditory rhythm on temporal stride parameters and EMG patterns in normal and hemiparetic gait. *Neurology*, 42, 208. [Abstract]

Thaut, M.H. & McIntosh, G.C. 1999. Music therapy and mobility training with the elderly: a review of current research. *Care Management Journals*, 1, 71-74.

Thaut, M.H. & McIntosh, G.C. 2006. Rhythmic auditory training in sensorimotor rehabilitation of people with Parkinson's Disease. *Neurorehabilitation & Neural Repair*, 20, 81.

Thaut, M.H., McIntosh, G.C., & Hoemberg, V. 1999. Mechanisms of rhythmic motor facilitation in hemiparetic stroke rehabilitation. *Journal of Neural Recovery & Repair*, 13, 30-31. [Abstract]

Thaut, M.H., McIntosh, G.C., Prassas, S.G., & Rice, R.R. 1992. Effects of auditory rhythmic pacing on normal gait and gait in stroke, cerebellar disorder and transverse myelitis. In M. Woollacott & F. Horak (Eds.), *Posture and Gait: Control Mechanisms*, Vol. 2 (pp. 437-440). Eugene, OR: University of Oregon Books.

Thaut, M.H., McIntosh, G.C., Prassas, S.G., & Rice, R.R. 1993. The effect of auditory rhythmic cuing on stride and EMG patterns in hemiparetic gait of stroke patients. *Journal of Neurologic Rehabilitation*, 7, 9-16.

Thaut, M.H., McIntosh, G.C., & Rice, R.R. 1997. Rhythmic facilitation of gait training in hemiparetic stroke rehabilitation. *Journal of Neurological Sciences*, 151, 207-212.

Thaut, M.H., McIntosh, G.C., Rice, R.R., & Prassas, S.G. 1993. The effect of auditory rhythmic cuing on stride and EMG patterns in hemiparetic gait of stroke patients. *Journal of Neurologic Rehabilitation*, 7, 9-16.

Thaut, M.H., McIntosh, G.C., Rice, R.R., & Miller, R.A. 1995. Rhythmic auditory-motor

training in gait rehabilitation with stroke patients. *Journal of Stroke and Cerebrovascular Disease*, 5, 100-101. [Abstract]

Thaut, M.H., McIntosh, G.C., Rice, R.R., Miller, R.A., Rathbun, J., & Brault, J.M. 1996. Rhythmic auditory stimulation in gait training with Parkinson's disease patients. *Movement Disorders*, 11, 193-200.

Thaut, M.H., McIntosh, G.C., Rice, R.R., & Prassas, S.G. 1993. Effect of auditory rhythmic cuing on temporal stride parameters and EMG patterns in hemiparetic gait of stroke patients. *Journal of Neurologic Rehabilitation*, 7, 9-16.

Thaut, M.H., McIntosh, G.C., Rice, R.R., Miller, R.A., Rathbun, J., & Brault, J.M. 1996. Rhythmic auditory stimulation in gait training of Parkinson's disease patients. *Movement Disorders*, 11, 193-200.

Thaut, M.H., McIntosh, K.H., McIntosh, G.C., & Hoemberg, V. 2001. Auditory rhythmicity enhances movement and speech motor control in patients with Parkinson's disease. *Functional Neurology*, 16, 163-172.

Thaut, M.H., Miller, R.A., Mezza, C.M., Rice, R.R., & McIntosh, G.C. 1995. Synchronization effects of auditory rhythm on gait healthy elderly and Parkinsonian patients on and off medication. *Proceedings Society for Neuroscience*, 819.1. [Abstract]

Thaut, M.H., Miltner, R., & Hoemberg, V. 1996. Rhythmisch-Akustische Stimulation in der Gangrehabilitation: Zusammenfassung bisheriger Befunde und Hinweise zur praktischen Durchfuehrung. *Neurologie & Rehabilitation*, 2, 81-86. [In German]

Thaut, M.H., Miltner, R., Lange, H.L., Hurt, C.P., & Hoemberg, V. 1997. Geschwindigkeitsmodulation und rhythmisch akustische Synchronisation des Ganges bei Patienten mit Morbus Huntington. *Neurologie & Rehabilitation*, 4 (suppl), 6. [Abstract]

Thaut, M.H., Miltner, R., Lange, H.L., Hurt, C.P., & Hoemberg, V. 1998. Gait velocity modulation with and without rhythmic facilitation in Huntington's disease patients. *Movement Disorders*, 14, 808-819.

Thaut, M.H., Miltner, R., Lange, H.L., Hurt, C.P., & Hoemberg, V. 1999. Velocity modulation and rhythmic synchronization of gait in Huntington's disease. *Movement Disorders*, 14 (5), 808-819.

Thaut, M.H., Nickel, A., & Hoemberg, V. 2003. Neurologische Musiktherapie: Ueberblick zum wissenschaftlichen Hintergrund und zur klinischen Methodik. *Musiktherapeutische Umschau*, In Press. [In German]

Thaut, M.H., Nickel, A., Kenyon, G.P., Meissner, N., & McIntosh, G.C. 2005. Rhythmic auditory stimulation (RAS) for gait training in hemiparetic stroke rehabilitation: An international multicenter study. *Proceedings Society for Neuroscience*, 756.6

Thaut, M.H., Rice, R.R., McIntosh, G.C., & Prassas, S.G. 1993. The effect of auditory rhythmic cuing on stride and EMG patterns in hemiparetic gait of stroke patients. *Physical Therapy*, 73, 107. [Abstract]

Thaut, M.H., Schicks, W., McIntosh, G.C., & Hoemberg, V. 2002. The role of motor imagery and temporal cuing in hemiparetic arm rehabilitation. *Journal of Neural Recovery & Repair*, In Press. [Abstract]

Thaut, M.H., Schleifers, S., & Davis, W.B. 1992. Changes in EMG patterns under the influence of auditory rhythm. In R. Spintge & R. Doh (Eds.), *Music Medicine* (pp. 80-101). St. Louis, MO: MMB Music.

Thaut, M.H., Ueno, K., Hurt, C.P., & Hoemberg, V. 1999. Bilateral limb entrainment and rhythmic synchronization in paretic arm movements of stroke patients. *Proceedings Society for Neuroscience*, 365.6. [Abstract]

Thaut, M.H., Schicks, W., McIntosh, G.C., & Hoemberg, V. (2002). The role of motor imagery and temporal cuing in hemiparetic arm rehabilitation. *Neurorehabilitation and Neural Repair*, 16:115.

Thaut, M.H., Leins, A.K., Rice, R.R., Argstatter, H., Kenyon, G.P., McIntosh, G.C., Bolay, H.V., Fetter, M. 2007. Rhythmic auditory stimulation improves gait more than NDT/Bobath Training in near-ambulatory patients early poststroke: A single- blind, randomized trial. *Neurorehabilitation and neural repair*, XX, 1-5.

Thompson, A.B., Arnold, J.C., & Murphy, S.E. 1990. Music therapy assessment of the cerebrovascular accident patient. *Music Therapy Perspectives*, 20, 1471-1485.

Trombetti A, Hars M, Herrmann FR, Kressig RW, Ferrari S, Rizzoli R. 2011. Effect of music-based multitask training on gait, balance, and fall risk in elderly people: a randomized controlled trial. *Arch Intern Med*. 171(6),525-33.

Van Wegen, W., Nieuwboer, A., Rochester, L., Jones, D., & Kwakkel, G. (2006). The effect of cueing therapy as measured by activity monitoring in the RESCUE trial. *Proceedings of the 4th World Congress for Neurorehabilitation*, P1-061.

Van Wegen, E., de Goede, C., Lim, I., Rietberg, M., Nieuwboer, A., Willems, A., Jones, D., Rochester, L., Hetherington, V., Berendse, H., Zijlmans, J., Wolters, E., & Kwakkel, G. 2006. The effect of rhythmic somatosensory cueing on gait in patients with Parkinson's disease. *Journal of the Neurological Sciences*, 248 (1-2), 210-4.

Van Wegen, E., Lim, I., de Goede, C., Nieuwboer, A., Willems, A., Jones, D., Roehster, L., Hetherington, V., Berendse, H., Zijlmans, J., Wolters, E., & Kwakkel, G. 2006. The effects of visual rhythms and optic flow on stride patterns of patients with Parkinson's disease. *Parkinsonism & Related Disorders*, 12(1), 21-7.

Willems, A.M., Nieuwboer, A., Chavret, F., Desloovere, K., Dom, R., Rochester, L., Jones, D., Kwakkel, G., & Van Wegen, E. 2006. The use of rhythmic auditory cues to influence gait in patients with Parkinson's disease, the differential effect for freezers and non-freezers, an explorative study. *Disability & Rehabilitation*, 28(11), 721-8.

Willems, A. 2006a. The effects of guideline-based cueing therapy on gait-related mobility in Parkinson's disease patients: The RESCUE-project. *Proceedings of the 4th World Congress for Neurorehabilitation*, F1B-2.

Willems, A. 2006b. Physiotherapy guidelines on the use of cueing in Parkinson's disease. Proceedings of the 4th World Congress for Neurorehabilitation, F1B-4.

Whitall, J., McCombe, Waller, S., Silver, K.H., & Macko, R.F. 2000. Repetitive bilateral arm training with rhythmic auditory cueing improves motor function in chronic hemiparetic stroke. *Stroke*, 31 (10), 2390-2395.

Yasuhara, A., Sugiyama, Y. 2001. Music therapy for children with Rett Syndrome. *Brain and Development*, 23, 1:S82-84.