

Bibliography

Sensorimotor

Rhythmic Auditory Stimulation (RAS)

- Alashram, A.R., Annino, G., & Mercuri, N. B. (2019). Rhythmic auditory stimulation in gait rehabilitation for traumatic brain and spinal cord injury. *Journal of Clinical Neuroscience*, *69*, 287-288.
doi: 10.1016/j.jocn.2019.08.080
- Arias, P., Cudeiro, J. (2008). Effects of rhythmic sensory stimulation (auditory, visual) on gait in Parkinson's disease patients. *Exp Brain Res*. 186 (4), 589-601.
doi: 10.1007/s00221-007-1263-y
- Baker, K., Rochester, L., & Nieuwboer, A. (2008). The effect of cues on gait variability—Reducing the attentional cost of walking in people with Parkinson's disease. *Parkinsonism & related disorders*, *14*(4), 314-320.
doi: 10.1016/j.parkreldis.2007.09.008
- Baram Y, Lenger R. (2012). Gait improvement in patients with cerebral palsy by visual and auditory feedback. *Neuromodulation*. 15(1),48-52.
doi: 10.1111/j.1525-1403.2011.00412.x
- Baram, Y. & Miller, A. (2007). Auditory feedback control for improvement of gait in patients with multiple sclerosis. *Neurological Sciences*, *254*, 90-94.
doi: 10.1016/j.jns.2007.01.003
- Benoit, C. E., Dalla Bella, S., Farrugia, N., Obrig, H., Mainka, S., & Kotz, S. A. (2014). Musically cued gait-training improves both perceptual and motor timing in Parkinson's disease. *Frontiers in human neuroscience*, *8*, 494.
doi: 10.3389/fnhum.2014.00494
- Bharathi, G., Jayaramayya, K., Balasubramanian, V., & Vellingiri, B. (2019). The potential role of rhythmic entrainment and music therapy intervention for individuals with autism spectrum disorders. *Journal of exercise rehabilitation*, *15*(2), 180–186.
doi:10.12965/jer.1836578.289
- Blecher, T., Tal, I., & Ben-Shachar, M. (2016). White matter microstructural properties correlate with sensorimotor synchronization abilities. *NeuroImage* *138*, 1–12.
doi: 10.1016/j.neuroimage.2016.05.022
- Bryant M.S., Rintala D.H., Lai E.C., Protas E.J. (2009). An evaluation of self-administration of auditory cueing to improve gait in people with Parkinson's disease. *Clinical Rehabilitation*. 23(12), 1078-1085.
doi: 10.1177/0269215509337465
- Buard, I., Dewispelaere, W. B., Teale, P., Rojas, D. C., Kronberg, E., Thaut, M. H., Kluger, B. M. (2019). Auditory entrainment of motor responses in older adults with and without Parkinson's disease: An MEG study. *Neuroscience Letters*, *708*.
doi.org/10.1016/j.neulet.2019.134331
- Bukowska, A. A., Krężałek, P., Mirek, E., Bujas, P., & Marchewka, A. (2015). Neurologic music therapy training for mobility and stability rehabilitation with Parkinson's disease—A pilot study. *Frontiers in human neuroscience*, *9*.
doi: 10.3389/fnhum.2015.00710
- Cameron, D. J., Pickett, K. A., Earhart, G. M., & Grahn, J. A. (2016). The effect of dopaminergic medication on beat-based auditory timing in Parkinson's disease. *Frontiers in Neurology* *7*, 1–8.
doi: 10.3389/fneur.2016.00019
- Carver, F. W., Fuchs, A., Jantzen, K. J., & Kelso, J. (2002). Spatiotemporal analysis of the neuromagnetic response to rhythmic auditory stimulation: Rate dependence and transient to steady-state transition. *Clinical Neurophysiology*, *113*(12), 1921-1931.
doi: 10.1016/s1388-2457(02)00299-7
- Cha, Y., Kim, Y., Hwang, S., & Chung, Y. (2014). Intensive gait training with rhythmic auditory stimulation in individuals with chronic hemiparetic stroke: A pilot randomized controlled study. *Neurorehabilitation* *35*(4), 681–688.
doi: 10.3233/nre-141182
- Clair AA, O'Konski M. (2006). The effect of rhythmic auditory stimulation (RAS) on gait characteristics of cadence, velocity, and stride length in persons with late stage dementia. *J Music Ther*. 43(2), 154-63.

- doi: 10.1093/jmt/43.2.154
- Conklyn, D., Stough, D., Novak, E., Paczak, S., Chemali, K., & Bethoux, F. (2010). A home-based walking program using rhythmic auditory stimulation improves gait performance in patients with multiple sclerosis: A pilot study. *Neurorehabilitation And Neural Repair*, 24(9), 835-842.
doi: 10.1093/jmt/43.2.154
- Cubo E., Leurgans S., Goetz C.G. (2004). Short-term and practice effects of metronome pacing in Parkinson's disease patients with gait freezing while in the 'on' state: randomized single blind evaluation. *Parkinsonism & Related Disorders*. 10(8), 507-510.
doi: 10.1016/j.parkreldis.2004.05.001
- de Bruin N., Doan J.B., Turnbull G., Suchowersky O., Bonfield S., Hu B., Brown L.A. (2010). Walking with music is a safe and viable tool for gait training in Parkinson's disease: the effect of a 13-week feasibility study on single and dual task walking. *Parkinsons Disorders*. 483530.
doi: 10.4061/2010/483530
- de Dreu M.J., van der Wilk A.S., Poppe E., Kwakkel G., van Wegen E.E. (2012). Rehabilitation, exercise therapy and music in patients with Parkinson's disease: a meta-analysis of the effects of music-based movement therapy on walking ability, balance and quality of life. *Parkinsonism & Related Disorders*. 18, S114-119.
doi: 10.1016/s1353-8020(11)70036-0
- de l'Etoile SK. (2008). The effect of rhythmic auditory stimulation on the gait parameters of patients with incomplete spinal cord injury: an exploratory pilot study. *Int J Rehabil Res*, 31(2),155-157.
doi: 10.1097/mrr.0b013e3282fc0f44
- Del Olmo, Aria P., Furio, M., Pozo, M., & Cudeiro, J. (2006). Evaluation of the effect of training using auditory stimulation on rhythmic movement in Parkinsonian patients—a combined motor and [18F]-FDG PET study. *Parkinsonism & Related Disorders*, 12(3), 155-164.
doi: 10.1016/j.parkreldis.2005.11.002
- Dozza, M., Chiari, L., Hlavacka, F., Cappello, A., & Horak, F.B. (2006). Effects of linear versus sigmoid coding of visual or audio biofeedback for the control of upright stance. *IEEE Transactions on Neural Systems & Rehabilitation Engineering*, 14(4), 505-512.
doi: 10.1109/tnsre.2006.886732
- Elsner, B., Schöler, A., Kon, T., & Mehrholz, J. (2019). Walking with rhythmic auditory stimulation in chronic patients after stroke: A pilot randomized controlled trial. *Physiother Res Int*.
doi: 10.1002/pri.1800
- Erra, C., Mileti, I., Germanotta, M., Petracca, M., Imbimbo, I., De Biase, A., ... & Padua, L. (2019). Immediate effects of rhythmic auditory stimulation on gait kinematics in Parkinson's disease ON/OFF medication. *Clinical Neurophysiology*, 130(10), 1789-1797.
doi: 10.1016/j.clinph.2019.07.013
- Fernández-del-Olmo, M., & Cudeiro Mazaira, F. J. (2003). A simple procedure using auditory stimuli to improve movement in Parkinson's disease: a pilot study.
[link:https://www.researchgate.net/publication/8258830_A_simple_procedure_using_auditory_stimuli_to_improve_movement_in_Parkinson's_disease_A_pilot_study](https://www.researchgate.net/publication/8258830_A_simple_procedure_using_auditory_stimuli_to_improve_movement_in_Parkinson's_disease_A_pilot_study)
- Ford, M., Wagenaar, R., Newell, K. (2007). The effects of auditory rhythms and instruction on walking patterns in individuals post stroke. *Gait and Posture*, 26, 150-155.
doi: 10.1016/j.gaitpost.2006.08.007
- Ford, M. P., Malone, L. A., Nyikos, I., Yelisetty, R., & Bickel, C. S. (2010). Gait training with progressive external auditory cueing in persons with Parkinson's disease. *Archives of physical medicine and rehabilitation*, 91(8), 1255-1261.
doi: 10.1016/j.apmr.2010.04.012
- Frazzitta, G, Maestri, R, Uccellini, D, Bertotti, G, Abelli, P. (2009). Rehabilitation treatment of gait in patients with Parkinson's disease with freezing: a comparison between two physical therapy protocols using visual and auditory cues with or without treadmill training. *Movement Disorders*. 24(8), 1139-43.
doi: 10.1002/mds.22491
- Freedland, R.L., Festa, C., Sealy, M., McBean, A., Elghazaly, P., Capan, A., Brozycki, L., Nelson, A.J., & Rothman, J. (2002). The effects of pulsed auditory stimulation on various gait measurements in persons with Parkinson's disease. *Neurorehabilitation*, 17 (1), 81-87.
doi: 10.3233/nre-2002-17110
- Freeman, J.S., Cody, F.W., & Schady, W. (1993). The influence of external timing cues upon the rhythm of voluntary movements in Parkinson's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*, 56, 1078-1084.
doi: 10.1136/jnnp.56.10.1078

- Georgiou, N., Iansek, R., Bradshaw, J. L., Phillips, J. G., Mattingley, J. B., & Bradshaw, J. A. (1993). An evaluation of the role of internal cues in the pathogenesis of parkinsonian hypokinesia. *Brain*, *116*(6), 1575-1587.
doi: 10.1093/brain/116.6.1575
- Ghai, S., Ghai, I., & Effenberg, A. O. (2017). Effect of rhythmic auditory cueing on gait in cerebral palsy: A systematic review and meta-analysis. *Neuropsychiatric Disease and Treatment* *14*, 43–59.
doi: 10.2147/ndt.s148053
- Ghai, S., Ghai, I., & Effenberg, A. O. (2018). Effect of rhythmic auditory cueing on aging gait: A systematic review and meta-analysis. *Aging and Disease* *9*(5), 901–923.
doi: 10.14336/ad.2017.1031
- Ghai, S., Ghai, I., Schmitz, G., & Effenberg, A. O. (2018). Effect of rhythmic auditory cueing on Parkinsonian gait: A systematic review and meta-analysis. *Scientific Reports* *8*, 506.
doi: 10.1038/s41598-017-16232-5
- Goldstroom Y, Knorr G, Goldstroom I. (2010). Rhythmic exercises in rehabilitation of TBI patients: a case report. *J Bodyw Mov Ther.* *14*(4), 336-45.
doi: 10.1016/j.jbmt.2009.06.002
- Hausdorff, J. M., Lowenthal, J., Herman, T., Gruendlinger, L., Peretz, C., & Giladi, N. (2007). Rhythmic auditory stimulation modulates gait variability in Parkinson's disease. *European Journal of Neuroscience*, *26*(8), 2369-2375.
doi: 10.1111/j.1460-9568.2007.05810.x
- Hesse, S. & Werner, C., & Bardeleben, A. (2004). Electromechanical gait training with functional electrical stimulation: Case studies in spinal cord injury. *Spinal Cord*, *42*, 346-52.
doi: 10.1038/sj.sc.3101595
- Hesse, S. & Werner, C. (2003). Post-stroke motor dysfunction and spasticity: Novel pharmacological and physical treatment strategies. *CNS Drugs*, *17*, 1070-1093.
doi: 10.2165/00023210-200317150-00004
- Hove, M. et al. (2012). Interactive rhythmic auditory stimulation reinstates natural 1/f timing in gait of Parkinson's patients. *PLoS ONE*, *7*(3).
doi: 10.1371/journal.pone.0032600
- Howe, T. E., Lövgreen, B., Cody, F. W., Ashton, V. J., & Oldham, J. A. (2003). Auditory cues can modify the gait of persons with early-stage Parkinson's disease: a method for enhancing parkinsonian walking performance? *Clinical Rehabilitation*, *17*(4), 363-367.
doi: 10.1191/0269215503cr621oa
- Hummelsheim, H. (1999). Rationales for improving motor function. *Current opinion in Neurology*, *12*(6), 697-701.
doi: 10.1097/00019052-199912000-00007
- Hurt, C.P., Rice, R.R., McIntosh, G.C., & Thaut, M.H. (1998). Rhythmic auditory stimulation in gait training for patients with traumatic brain injury. *Journal of Music Therapy*, *35*, 228-241.
doi: 10.1093/jmt/35.4.228
- Jeffery, D.R. & Good, D.C. (1995). Rehabilitation of the stroke patient. *Current Opinion in Neurology*, *8*, 62-68.
doi: 10.1097/00019052-199502000-00010
- Jeong, S., Kim, M.T. (2007). Effects of a theory-driven music and movement program for stroke survivors in a community setting. *Applied Nursing Research*, *20*(3), 125-31.
doi: 10.1016/j.apnr.2007.04.005
- Kadivar, Z., Corcos, D. M., Foto, J., & Hondzinski, J. M. (2011). Effect of step training and rhythmic auditory stimulation on functional performance in Parkinson patients. *Neurorehabilitation and neural repair*, *25*(7), 626-635.
doi: 10.1177/1545968311401627
- Kaipust, J. P., McGrath, D., Mukherjee, M., & Stergiou, N. (2013). Gait variability is altered in older adults when listening to auditory stimuli with differing temporal structures. *Annals of biomedical engineering*, *41*(8), 1595-1603.
doi: 10.1007/s10439-012-0654-9
- Kelly, G., & Pool, J. (2019). Rhythmic auditory stimulation in gait rehabilitation for children and youth following acquired brain injury. *International Journal of Therapy and Rehabilitation*, *26*(6), 12-12.
doi: 10.12968/ijtr.2019.26.6.12
- Kim J.S., Oh D.W., Kim S.Y., Choi J.D. (2011). Visual and kinesthetic locomotor imagery training integrated with auditory step rhythm for walking performance of patients with chronic stroke. *Clin Rehabil.* *25*(2),134-45.
doi: 10.1177/0269215510380822

- Kim, S. J., Kwak, E. E., Park, E. S., & Cho, S. R. (2012). Differential effects of rhythmic auditory stimulation and neurodevelopmental treatment/Bobath on gait patterns in adults with cerebral palsy: a randomized controlled trial. *Clinical rehabilitation*, 26(10), 904-914.
doi: 10.1177/0269215511434648
- Kim, S. J., Kwak, E. E., Park, E. S., Lee, D. S., Kim, K. J., Song, J. E., & Cho, S. R. (2011). Changes in gait patterns with rhythmic auditory stimulation in adults with cerebral palsy. *NeuroRehabilitation*, 29(3), 233-241.
doi: 10.3233/nre-2011-0698
- Kim, J. S., & Oh, D. W. (2012). Home-based auditory stimulation training for gait rehabilitation of chronic stroke patients. *Journal of Physical Therapy Science*, 24(8), 775-777.
doi: 10.1589/jpts.24.775
- Koshimori, Y., Strafella, A., Valli, M., Sharma, V., Cho, S. S., Houle, S., & Thaut, M. H. (2018). Motor synchronization to rhythmic auditory stimulation (RAS) attenuates dopamine response in the ventral striatum in young healthy adults. *Proceedings of the Society for Neuroscience* 670, 15.
doi: 10.3389/fnins.2019.00106
- Kwak, E.E. (2007). Effect of rhythmic auditory stimulation on gait performance in children with spastic cerebral palsy. *Journal of Music Therapy*, 44, 198-216.
doi: 10.1093/jmt/44.3.198
- Ledger, s, Galvin, R, Lynch, D, Stokes, EK. (2008). A randomized controlled trial evaluating the effect of an individual auditory cueing device on freezing and gait speed in people with Parkinson's disease. *BMC Neurology*. 8, 46.
doi: 10.1186/1471-2377-8-46
- Lei, J., Conradi, N., Abel, C., Frisch, S., Brodski-Guerniero, A., Hildner, M., ... & Schmidt-Kassow, M. (2019). Cognitive effects of rhythmic auditory stimulation in Parkinson's disease: A P300 study. *Brain Research*, 1716, 70-79.
doi: 10.1016/j.brainres.2018.05.016
- de L'Etoile, S.K. (2008). The effect of rhythmic auditory stimulation on the gait parameters of patients with incomplete spinal cord injury: An exploratory pilot study. *International Journal of Rehabilitation Research*, 31, 155-157.
doi: 10.1097/mrr.0b013e3282fc0f44
- Lim, I., van Wegen, E., de Goede, C., Deutekom, M., Nieuwboer, A., Willems, A., ... & Kwakkel, G. (2005). Effects of external rhythmical cueing on gait in patients with Parkinson's disease: a systematic review. *Clinical rehabilitation*, 19(7), 695-713.
doi: 10.1191/0269215505cr906oa
- Lohnes, CA, Earhart, GM. (2011). The impact of attentional, auditory, and combined cues on walking during single and cognitive dual tasks in Parkinson disease. *Gait & Posture*. 33 (3), 478-83.
doi: 10.1016/j.gaitpost.2010.12.029
- Luft, A.R., McCombe-Waller, S., Whittall, J., Forrester, L.W., Macko, R., Sorkin, J.D., Schulz, J.B., Goldberg, A.P., & Hanley, D.F. (2004). Repetitive bilateral arm training and motor cortex activation in chronic stroke: A randomized controlled trial. *JAMA*, 292 (15), 1853-1861.
doi: 10.1001/jama.292.15.1853
- Ma, H.I., Hwang, W.J., & Lin, K.C. (2009). The effects of two different auditory stimuli on functional arm movement in persons with Parkinson's disease: a dual-task paradigm. *Clinical Rehab*, 23, 229-237.
doi: 10.1177/0269215508098896
- Malcolm, M. P., Massie, C., & Thaut, M. (2009). Rhythmic auditory-motor entrainment improves hemiparetic arm kinematics during reaching movements: a pilot study. *Topics in stroke rehabilitation*, 16(1), 69-79.
doi: 10.1310/tsr1601-69
- Malherbe, V., Brenière, Y., & Bril, B. (1992). How do cerebral palsied children with hemiplegia control their gait. *Posture and control mechanisms*, 2.
- Mandel, A.R., Nymark, J.R., Balmer, S.J., Grinnell, D.M., & O'Riain, M.D. (1990). Electromyographic feedback versus rhythmic positional biofeedback in computerized gait retraining with stroke patients. *Archives of Physical and Medical Rehabilitation*, 71, 649-654.
link: <https://www.ncbi.nlm.nih.gov/pubmed/2375668>
- Mauritz, K.H. (2002). Gait training in hemiplegia. *European Journal of Neurology*, 9(1)1, 23-29, discussion 53-61.
doi: 10.1046/j.1468-1331.2002.0090s1023.x
- McIntosh, G.C., Brown, S.H., Rice, R.R., & Thaut, M.H. (1997). Rhythmic auditory-motor facilitation of gait patterns in patients with Parkinson's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*, 62,

122-126.

doi: 10.1136/jnnp.62.1.22

- McIntosh, G.C., Rice, R.R., Hurt, C.P., & Thaut, M.H. (1998). Long-term training effects of rhythmic auditory stimulation on gait in patients with Parkinson's disease. *Movement Disorders*, 13 (2), 212. [Abstract]
doi: 10.1093/jmt/35.4.228
- Miller, R.A., Thaut, M.H., McIntosh, G.C., Rice, R.R. (1996). Components of EMG symmetry and variability in Parkinsonian and healthy elderly gait. *Electroencephalography and Clinical Neurophysiology*, 101, 1-7.
doi: 10.1016/0013-4694(95)00209-x
- Mitoma H, Yoneyama M, Orimo S. (2010). 24-hour recording of parkinsonian gait using a portable gait rhythmogram. *Intern Med*. 49(22),2401-2408.
doi: 10.2169/internalmedicine.49.3511
- Morris, G.S., Suteerawattananon, M., Etnyre, B.R., Jankovic, J., & Protas, E.J. (2004). Effects of visual and auditory cues on gait in individuals with Parkinson's disease. *Journal of the Neurological Sciences*, 219, 63-69.
doi: 10.1016/j.jns.2003.12.007
- Munneke, M., Keus, S.H.J., Bredero-Cohen, A.B., Hendriks, H.J., & Bloem, B.R. (2006). Evidence-based clinical practice guideline for physical therapy in Parkinson's disease. *Proceedings of the 4th World Congress for Neurorehabilitation*, P1-058
doi: 10.1002/mds.21244
- Nanhoe-Mahabier W., Delval A., Snijders A.H., Weerdesteijn V., Overeem S., Bloem B.R. (2012). The possible price of auditory cueing: influence on obstacle avoidance in Parkinson's disease. *Movement Disorders*. 27(4), 574-578.
doi: 10.1002/mds.24935
- Nieuwboer, A., Kwakkel, G., Rochester, L., Jones, D., van Wegen E., Willems, A.M., Chavret, F., Hetherington, V., Baker, K., & Lim, I. (2007). Cueing training in the home improves gait-related mobility in Parkinson's disease: the RESCUE trial. *Journal of Neurology, Neurosurgery & Psychiatry*, 78(2), 134-40.
doi: 10.1136/jnnp.200x.097923
- Nombela, C., Hughes, L. E., Owen, A. M., & Grahn, J. A. (2013). Into the groove: can rhythm influence Parkinson's disease? *Neuroscience & Biobehavioral Reviews*, 37(10), 2564-2570.
doi: 10.1016/j.neubiorev.2013.08.003
- Pacchetti, C., Mancini, F., Aglieri, R., Fundarò, C., Martignoni, E., & Nappi, G. (2000). Active music therapy in Parkinson's disease: an integrative method for motor and emotional rehabilitation. *Psychosomatic medicine*, 62(3), 386-393.
doi: 10.1097/00006842-200005000-00012
- Paltsev Y.I. & Elnor, A.M. (1967). Change in the functional state of the segmental apparatus of the spinal cord under the influence of sound stimuli and its role in voluntary movement. *Biophysics*, 12, 1219-1226.
- Pau, M., Corona, F., Pili, R., Casula, C., Sors, F., Agostini, T., ... & Murgia, M. (2016). Effects of physical rehabilitation integrated with rhythmic auditory stimulation on spatio-temporal and kinematic parameters of gait in Parkinson's disease. *Frontiers in Neurology*, 7.
doi: 10.3389/fneur.2016.00126
- Pelton, T. A., Johannsen, L., Chen, H., & Wing, A. M. (2010). Hemiparetic stepping to the beat: asymmetric response to metronome phase shift during treadmill gait. *Neurorehabilitation and Neural Repair*, 24(5), 428-434.
doi: 10.1177/1545968309353608
- Pfeiffer, K., Clements, J., Smith, M., Gregoire, M., & Conti, C. (2019). Does rhythmic auditory stimulation compared to no rhythmic auditory stimulation improve patient's static and dynamic standing balance post stroke? *Physical Therapy Reviews*, 24(5), 216-222.
doi: 10.1080/10833196.2019.1666220
- Picelli A, Camin M, Tinazzi M, Vangelista A, Cosentino A, Fiaschi A, Smania N. (2010). Three-dimensional motion analysis of the effects of auditory cueing on gait pattern in patients with Parkinson's disease: a preliminary investigation. *Neurol Sci*. 31(4),423-30.
doi: 10.1007/s10072-010-0228-2
- Prassas, S., Thaut, M., McIntosh, G., & Rice, R. (1997). Effect of auditory rhythmic cuing on gait kinematic parameters of stroke patients. *Gait & Posture*, 6(3), 218-223.
doi: 10.1016/s0966-6362(97)00010-6
- Rao, S. M., Mayer, A. R., & Harrington, D. L. (2001). The evolution of brain activation during temporal processing. *Nature neuroscience*, 4(3), 317-323. doi: 10.1038/85191

- Richards, C.L., Malouin, F., Bedard, P.J., & Cioni, M. (1992). Changes induced by L-Dopa and sensory cues on the gait of parkinsonian patients. In M. Woollacott and F. Horak (Eds.), *Posture and Control Mechanisms* (Vol. 2) (pp. 126-129). Eugene, OR: University of Oregon Books.
- Rizzonelli, M., Kim, J. H., Gladow, T., & Mainka, S. (2017). Musical stimulation with feedback in gait training for Parkinson's disease. *Psychomusicology: Music, Mind, and Brain* 27, 213–218.
doi: 10.1037/pmu0000182
- Rochester, L., Baker, K., Nieuwboer, A., & Burn, D. (2011). Targeting dopa-sensitive and dopa-resistant gait dysfunction in Parkinson's disease: Selective responses to internal and external cues. *Movement Disorders*, 26(3), 430-435.
doi: 10.1002/mds.23450
- Roerdink, M. et al. (2011). Walking to the beat of different drums: Practical implications for the use of acoustic rhythms in gait rehabilitation. *Gait and Posture*, 33(1), 690-694.
doi: 10.1016/j.gaitpost.2011.03.001
- Roerdink, M., Lamoth, C.J.C., van Kordelaar, J., Elich, P., Konijnenbelt, M., Kwakkel, G. & Beek, P.J. (2009). Rhythm perturbations in acoustically paced treadmill walking after stroke. *Neurorehabilitation and Neural Repair*, 23, 668-678.
doi: 10.1016/j.gaitpost.2009.08.193
- Roerdink, M., Lamoth, C.J.C., Kwakkel, G., van Wieringen, P.C.W. & Beek, P.J. (2007). Gait coordination after stroke: Benefits of acoustically paced treadmill walking. *Physical Therapy*, 87, 1009-1022.
doi: 10.2522/ptj.20050394
- Rossignol, S. & Melvill Jones, G. (1976). Audiospinal influences in man studied by the H-reflex and its possible role in rhythmic movement synchronized to sound. *Electroencephalography & Clinical Neurophysiology*, 41 (1), 83-92.
doi: 10.1016/0013-4694(76)90217-0
- Safranek, M.G., Koshland, G.F., & Raymond, G. (1982). The influence of auditory rhythm on muscle activity. *Physical Therapy*, 2, 161-168.
doi: 10.1093/ptj/62.2.161
- Satoh, M. & Kuzuhara, S. (2008). Training in mental singing while walking improves gait disturbance in Parkinson's disease patients. *European Neurology*, 60, 237-243.
doi: 10.1159/000151699
- Schauer, M. & Mauritz, K.H. (2003). Musical motor feedback (MMF) in walking hemiparetic stroke patients: Randomized trials of gait improvement. *Clinical Rehabilitation*, 17, 713-722.
doi: 10.1191/0269215503cr668oa
- Secoli, R. et al. (2011). Effect of visual distraction and auditory feedback on patient effort during robot-assisted movement training after stroke. *Journal of Neuroengineering and Rehabilitation*, 8(21).
doi: 10.1186/1743-0003-8-21
- Seebacher, B., Kuisma, R., Glynn, A., & Berger, T. (2016). The effect of rhythmic-cued motor imagery on walking, fatigue and quality of life in people with multiple sclerosis: A randomised controlled trial. *Multiple Sclerosis Journal* 23(2), 286–296.
doi: 10.1177/1352458516644058
- Shahraki, M., Sohrabi, M., Torbati, H. T., Nikkhah, K., & Naeimi Kia, M. (2017). Effect of rhythmic auditory stimulation on gait kinematic parameters of patients with multiple sclerosis. *Journal of Medicine and Life* 10(1), 33–37.
link: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5304368/>
- Shin, Y. K., Chong, H. J., Kim, S. J., & Cho, S. R. (2015). Effect of rhythmic auditory stimulation on hemiplegic gait patterns. *Yonsei medical journal*, 56(6), 1703-1713.
doi: 10.3349/ymj.2015.56.6.1703
- Song, G. B., & Ryu, H. J. (2016). Effects of gait training with rhythmic auditory stimulation on gait ability in stroke patients. *Journal of Physical Therapy Science* 28(5), 1403– 1406.
doi: 10.1589/jpts.28.1403
- Song, J. H., Zhou, P. Y., Cao, Z. H., Ding, Z. G., Chen, H. X., & Zhang, G. B. (2015). Rhythmic auditory stimulation with visual stimuli on motor and balance function of patients with Parkinson's disease. *European Review for Medical and Pharmacological Sciences* 19(11), 2001–2007.
link: [khttps://www.researchgate.net/publication/279628910_Rhythmic_auditory_stimulation_with_visual_stimuli_on_motor_and_balance_function_of_patients_with_Parkinson's_disease](https://www.researchgate.net/publication/279628910_Rhythmic_auditory_stimulation_with_visual_stimuli_on_motor_and_balance_function_of_patients_with_Parkinson's_disease)
- Spaulding, S. J., Barber, B., Colby, M., Cormack, B., Mick, T., & Jenkins, M. E. (2013). Cueing and gait improvement among people with Parkinson's disease: a meta-analysis. *Archives of physical medicine and rehabilitation*, 94(3), 562-570.
doi: 10.1016/j.apmr.2012.10.026

- Stephan, K.M., Thaut, M.H., Wunderlich, G., Schicks, W., Tian, B., Tellmann, L., Schmitz, T., Herzog, H., McIntosh, G.C., Seitz, R.J., Hoemberg, V. (2002). Conscious and subconscious sensorimotor synchronization: Prefrontal cortex and the influence of awareness. *Neuroimage*, 15 (2), 345-352. doi: 10.1006/nimg.2001.0929
- Stephan, K.M., Thaut, M.H., Wunderlich, G., Schicks, W., Tellmann, L., Herzog, H., McIntosh, G.C., Kraemer, M., Hoemberg, V., & Seitz, R.J. (2002). Cortico-cerebellar circuits and temporal adjustments of motor behavior. *Proceedings Society for Neuroscience*, 462.8. [Abstract]
- Styns, F., van Noorden, L., Moelants, D., & Leman, M. (2007). Walking on music. *Human movement science*, 26(5), 769-785. doi: 10.1016/j.humov.2007.07.007
- Suzuki, K., Niitsu, M., Kamo, T., Otake, S., & Nishida, Y. (2019). Effect of exercise with rhythmic auditory stimulation on muscle coordination and gait stability in patients with diabetic peripheral neuropathy: A randomized controlled trial. *Open Journal of Therapy and Rehabilitation*, 7, 79-91. doi: 10.4236/ojtr.2019.73005
- Teasell, R. W., Bhogal, S. K., Foley, N. C., & Speechley, M. R. (2003). Gait retraining post stroke. *Topics in stroke rehabilitation*, 10(2), 34-65. doi: 10.1310/udxe-mjff-53v2-eap0
- Thaut, C., & Stephan, K. M. (2019). Neurologic Music Therapy in Sensorimotor Rehabilitation. In M. H. Thaut & D. A. Hodges (Eds.), *The Oxford Handbook of Music and the Brain* (pp. 695-714). Oxford, England: Oxford University Press.
- Thaut, M.H., Hoemberg, V., Kenyon, G., & Hurt, C.P. (1998). Rhythmic entrainment of hemiparetic arm movements in stroke patients. *Proceedings Society for Neuroscience*, 653.7. [Abstract]
- Thaut, M.H., McIntosh, G.C., Prassas, S.G., & Rice, R.R. (1993). The effect of auditory rhythmic cuing on stride and EMG patterns in hemiparetic gait of stroke patients. *Journal of Neurologic Rehabilitation*, 7, 9-16. doi: 10.1177/136140969300700103
- Thaut, M. H., & Kenyon, G. P. (2003). Rapid motor adaptations to subliminal frequency shifts during syncopated rhythmic sensorimotor synchronization. *Human movement science*, 22(3), 321-338. doi: 10.1016/s0167-9457(03)00048-4
- Thaut, M.H., Kenyon, G.P., Hurt, C.P., McIntosh, G.C., & Hoemberg, V. (2002). Kinematic optimization of spatiotemporal patterns in paretic arm training with stroke patients. *Neuropsychologia*, 40, 1073-1081. doi: 10.1016/s0028-3932(01)00141-5
- Thaut, M.H., Lange, H., Miltner, R., Hurt, C.P., & Hoemberg, V. (1996). Rhythmic entrainment of gait patterns in Huntington's disease patients. *Proceedings Society for Neuroscience*, 727.6. [Abstract]
- Thaut, M. H., Leins, A. K., Rice, R. R., Argstatter, H., Kenyon, G. P., McIntosh, G. C., ... & Fetter, M. (2007). Rhythmic auditory stimulation improves gait more than NDT/Bobath training in near-ambulatory patients early poststroke: a single-blind, randomized trial. *Neurorehabilitation and neural repair*, 21(5), 455-459. doi: 10.1177/1545968307300523
- Thaut, M.H., McIntosh, G.C., & Rice, R.R. (1997). Rhythmic facilitation of gait training in hemiparetic stroke rehabilitation. *Journal of Neurological Sciences*, 151, 207-212. doi: 10.1016/s0022-510x(97)00146-9
- Thaut, M.H., McIntosh, K.H., McIntosh, G.C., & Hoemberg, V. (2001). Auditory rhythmicity enhances movement and speech motor control in patients with Parkinson's disease. *Functional Neurology*, 16, 163-172. link: <https://www.ncbi.nlm.nih.gov/pubmed/11495422>
- Thaut, M.H. & McIntosh, G.C. (2006). Rhythmic auditory training in sensorimotor rehabilitation of people with Parkinson's Disease. *Neurorehabilitation & Neural Repair*, 20, 81.
- Thaut, M.H., Miltner, R., Lange, H.L., Hurt, C.P., & Hoemberg, V. (1999). Velocity modulation and rhythmic synchronization of gait in Huntington's disease. *Movement Disorders*, 14 (5), 808-819. doi: 10.1002/1531-8257(199909)14:5<808::aid-mds1014>3.0.co;2-j
- Thaut, M.H., Nickel, A., Kenyon, G.P., Meissner, N., & McIntosh, G.C. (2005). Rhythmic auditory stimulation (RAS) for gait training in hemiparetic stroke rehabilitation: An international multicenter study. *Proceedings Society for Neuroscience*, 756.
- Thaut, M. H., Rice, R. R., Braun Janzen, T., Hurt-Thaut, C. P., & McIntosh, G. C. (2018). Rhythmic auditory stimulation for reduction of falls in Parkinson's disease: A randomized controlled study. *Clinical Rehabilitation*. doi:10.1177/0269215518788615. doi: 10.1177/0269215518788615
- Thaut, M.H., Schicks, W., McIntosh, G.C., & Hoemberg, V. (2002). The role of motor imagery and temporal cuing in hemiparetic arm rehabilitation. *Neurorehabilitation and Neural Repair*, 16, 115.

- Trombetti, A., Hars, M., Herrmann, F. R., Kressig, R. W., Ferrari, S., & Rizzoli, R. (2011). Effect of music-based multitask training on gait, balance, and fall risk in elderly people: a randomized controlled trial. *Archives of internal medicine*, *171*(6), 525-533.
doi: 10.1001/archinternmed.2010.446
- Uchitomi, H., Ota, L., Ogawa, K. I., Orimo, S., & Miyake, Y. (2013). Interactive rhythmic cue facilitates gait relearning in patients with Parkinson's disease. *PloS one*, *8*(9), e72176.
doi: 10.1371/journal.pone.0072176
- Van Wegen, E., de Goede, C., Lim, I., Rietberg, M., Nieuwboer, A., Willems, A., Jones, D., Rochester, L., Hetherington, V., Berendse, H., Zijlmans, J., Wolters, E., & Kwakkel, G. (2006). The effect of rhythmic somatosensory cueing on gait in patients with Parkinson's disease. *Journal of the Neurological Sciences*, *248* (1-2), 210-214.
doi: 10.1016/j.jns.2006.05.034
- Varsamis, P., Staikopoulos, K., & Kartasidou, L. (2012). Effect of Rhythmic Auditory Stimulation on Controlling Stepping Cadence of Individuals with Mental Retardation and Cerebral Palsy. *International Journal of Special Education*, *27*(3), 68-75.
link:https://www.researchgate.net/publication/236650155_Effect_of_rhythmic_auditory_stimulation_on_controlling_stepping_cadence_of_individuals_with_mental_retardation_and_cerebral_palsy
- Whitall, J., McCombe, Waller, S., Silver, K.H., & Macko, R.F. (2000). Repetitive bilateral arm training with rhythmic auditory cueing improves motor function in chronic hemiparetic stroke. *Stroke*, *31* (10), 2390-2395.
doi: 10.1161/01.str.31.10.2390
- Willems, A.M., Nieuwboer, A., Chavret, F., Desloovere, K., Dom, R., Rochester, L., Jones, D., Kwakkel, G., & Van Wegen, E. (2006). The use of rhythmic auditory cues to influence gait in patients with Parkinson's disease, the differential effect for freezers and non-freezers, an explorative study. *Disability & Rehabilitation*, *28*(11), 721-8.
doi: 10.1080/09638280500386569
- Willems, A. (2006a). The effects of guideline-based cueing therapy on gait-related mobility in Parkinson's disease patients: The RESCUE-project. *Proceedings of the 4th World Congress for Neurorehabilitation*, F1B-2.
doi: 10.1016/s0966-6362(05)80320-0
- Willems, A. (2006b). Physiotherapy guidelines on the use of cueing in Parkinson's disease. *Proceedings of the 4th World Congress for Neurorehabilitation*, F1B-4.
- Wittwer, J. E., Webster, K. E., & Hill, K. (2013). Rhythmic auditory cueing to improve walking in patients with neurological conditions other than Parkinson's disease—what is the evidence? *Disability and rehabilitation*, *35*(2), 164-176.
doi: 10.3109/09638288.2012.690495
- Wittwer, J. E., Webster, K. E., & Hill, K. (2013). Effect of rhythmic auditory cueing on gait in people with Alzheimer disease. *Archives of physical medicine and rehabilitation*, *94*(4), 718-724.
doi: 10.1016/j.apmr.2012.11.009
- Wright, R. L., Bevins, J. W., Pratt, D., Sackley, C. M., & Wing, A. M. (2016). Metronome cueing of walking reduces gait variability after a cerebellar stroke. *Frontiers in Neurology*, *7*.
doi: 10.3389/fneur.2016.00084
- Yoo, G. E., & Kim, S. J. (2016). Rhythmic auditory cueing in motor rehabilitation for stroke patients: Systematic review and meta-analysis. *Journal of Music Therapy* *53*(2), 149–177.
doi: 10.1093/jmt/thw003

Pattern Sensory Enhancement (PSE)

- Ackerley, S. J., Stinear, C. M., & Byblow, W. D. (2011). Promoting use-dependent plasticity with externally-paced training. *Clinical Neurophysiology*, *122*(12), 2462-2468.
doi: 10.1016/j.clinph.2011.05.011
- Bernardi NF, Aggujaro S, Caimmi M, Molteni F, Maravita A, Luzzatti C. (2009). A new approach to rhythm cueing of cognitive functions: the case of ideomotor apraxia. *Ann NY Acad Sci.* (1169),417-421.
doi: 10.1111/j.1749-6632.2009.04854.x
- Bernatzky, G., Bernatzky, P., Hesse, H.P., Staffen, W., & Ladurner, G. (2004). Stimulating music increases motor coordination in patients afflicted by Morbus Parkinson. *Neuroscience Letters*, *361*, 4-8.
doi: 10.1016/j.neulet.2003.12.022

- Brown, S. H., Thaut, M. H., Benjamin, J., & Cooke, J. D. (1993). Effects of rhythmic auditory cueing on temporal sequencing of complex arm movements. *Proceedings of the Society for Neuroscience*, 227 (2).
- Bukowska, A. A., Krężałek, P., Mirek, E., Bujas, P., & Marchewka, A. (2015). Neurologic music therapy training for mobility and stability rehabilitation with Parkinson's disease—A pilot study. *Frontiers in human neuroscience*, 9. doi: 10.3389/fnhum.2015.00710
- Buetefish, C., Hummelsheim, H., Denzler, P., & Mauritz, K.H. (1995). Repetitive training of isolated movements improves the outcome of motor rehabilitation of the centrally paretic hand. *Journal of Neurological Sciences*, 130, 59-68. doi: 10.1016/0022-510x(95)00003-k
- Del Olmo, Aria P., Furio, M., Pozo, M., & Cudeiro, J. (2006). Evaluation of the effect of training using auditory stimulation on rhythmic movement in Parkinsonian patients—a combined motor and [18F]-FDG PET study. *Parkinsonism & Related Disorders*, 12(3), 155-64. doi: 10.1016/j.parkreldis.2005.11.002
- Dozza, M., Chiari, L., Hlavacka, F., Cappello, A., & Horak, F.B. (2006). Effects of linear versus sigmoid coding of visual or audio biofeedback for the control of upright stance. *IEEE Transactions on Neural Systems & Rehabilitation Engineering*, 14(4), 505-12. doi: 10.1109/tnsre.2006.886732
- Dozza, M., Chiari, L., & Horak, F.B. (2005). Audio-biofeedback improves balance in patients with bilateral vestibular loss. *Archives of Physical Medicine and Rehabilitation*, 86 (7), 1401-3. doi: 10.1016/j.apmr.2004.12.036
- Effenberg, A. O., & Mechling, H. (1998). Bewegung hörbar machen—Warum? Zur Perspektive einer systematischen Umsetzung von Bewegung in Klänge. *psychologie und sport*, 1, 29-38.
- Freeman, J.S., Cody, F.W., & Schady, W. (1993). The influence of external timing cues upon the rhythm of voluntary movements in Parkinson's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*, 56, 1078-1084. doi: 10.1136/jnnp.56.10.1078
- Georgiou, N., Iansek, R., Bradshaw, J. L., Phillips, J. G., Mattingley, J. B., & Bradshaw, J. A. (1993). An evaluation of the role of internal cues in the pathogenesis of parkinsonian hypokinesia. *Brain*, 116(6), 1575-1587. doi: 10.1093/brain/116.6.1575
- Goldshtroum Y, Knorr G, Goldshtroum I. (2010). Rhythmic exercises in rehabilitation of TBI patients: a case report. *J Bodyw Mov Ther*. 14(4), 336-345. doi: 10.1016/j.jbmt.2009.06.002
- Jakubowski, K., Farrugia, N., & Stewart, L. (2016). Probing imagined tempo for music: Effects of motor engagement and musical experience. *Psychology of Music*, 44(6), 1274–1288. doi: 10.1177/0305735615625791
- Jeong, S., Kim, M.T. (2007). Effects of a theory-driven music and movement program for stroke survivors in a community setting. *Applied Nursing Research*, 20(3), 125-131. doi: 10.1016/j.apnr.2007.04.005
- Luft, A.R., McCombe-Waller, S., Whittall, J., Forrester, L.W., Macko, R., Sorkin, J.D., Schulz, J.B., Goldberg, A.P., & Hanley, D.F. (2004). Repetitive bilateral arm training and motor cortex activation in chronic stroke: A randomized controlled trial. *JAMA*, 292 (15), 1853-1861 doi: 10.1001/jama.292.15.1853
- Mak, M. (2006). Feed-forward audio-visual cues could enhance sit-to-stand in Parkinsonian patients. *Proceedings of the 4th World Congress for Neurorehabilitation*, F1B-7.
- Malcolm, M.P., Massie, C., Thaut, M.H. (2009). Rhythmic auditory- motor entrainment improves hemiparetic arm kinematics during reaching movements: A pilot study. *Top Stroke Rehabil*, 16, 69-79. doi: 10.1310/tsr1601-69
- McCombe, W.S. & Whittall, J. (2005). Hand dominance and side of stroke affect rehabilitation in chronic stroke. *Clinical Rehabilitation*, 19 (5), 544-551. doi: 10.1191/0269215505cr829oa
- McCombe Waller, S. & Whittall, J. (2006). Combining bilateral and distal arm training to promote arm and hand recovery in patients with chronic hemiparesis: A case report. *Proceedings of the 4th World Congress for Neurorehabilitation*, P2-080
- Pacchetti, C., Aglieri, R., Mancini, F., Martignoni, E., & Nappi, G. (1998). Active music therapy and Parkinson's disease: Methods. *Functional Neurology*, 13 (1), 57-67.
- Peng, Y. C., Lu, T. W., Wang, T. H., Chen, Y. L., Liao, H. F., Lin, K. H., & Tang, P. F. (2011). Immediate effects

- of therapeutic music on loaded sit-to-stand movement in children with spastic diplegia. *Gait & posture*, 33(2), 274-278.
doi: 10.1016/j.gaitpost.2010.11.020
- Robertson, S.D., Chua, R., Maraj, B.K., Kao, J.C., & Weeks, D.J. (2002a). Bimanual coordination dynamics in adults with downs syndrome. *Motor Control*, 6, 388-407.
doi: 10.1123/mcj.6.4.388
- Robertson, S.D., Van Gemmert, A.W., & Maraj, B.K. (2002b). Auditory information is beneficial for adults with downs syndrome in a continuous bimanual task. *Acta Psychologica*, 110, 213-229.
doi: 10.1016/s0001-6918(02)00034-3
- Sameiro-Barbosa, C. M., & Geiser, E. (2016). Sensory entrainment mechanisms in auditory perception: Neural synchronization cortico-striatal activation. *Frontiers in Neuroscience*.
doi: 10.3389/fnins.2016.00361
- Sandrini, G., Tassorelli, C., Balloni, L., Buscone, S., & Pacchetti, C. (2006). Efficacy of sensorial cues on gait rehabilitation in Parkinson's disease. *Proceedings of the 4th World Congress for Neurorehabilitation*, P1-053.
link: <https://journals.sagepub.com/doi/pdf/10.1177/1545968305284198>
- Senesac, Davis, and Richards. (2010). Generalization of a modified form of repetitive rhythmic bilateral training in stroke. *Human Movement Science*, 26(1), 137-148.
doi: 10.1016/j.humov.2009.05.004
- Simkins, M., Jacobs, A. B., & Rosen, J. (2013). Rhythmic affects on stroke-induced joint synergies across a range of speeds. *Experimental brain research*, 229(4), 517-524.
doi: 10.1007/s00221-013-3613-2
- Stephan, M. A., Brown, R., Lega, C., & Penhune, V. (2016). Melodic priming of motor sequence performance: The role of the dorsal premotor cortex. *Frontiers in Neuroscience*, 10, 210.
doi: 10.3389/fnins.2016.00210
- Stinear, J. W., & Byblow, W. D. (2004). Rhythmic bilateral movement training modulates corticomotor excitability and enhances upper limb motricity poststroke: a pilot study. *Journal of Clinical Neurophysiology*, 21(2), 124-131.
doi: 10.1097/00004691-200403000-00008
- Thaut, C., & Stephan, K. M. (2019). Neurologic Music Therapy in Sensorimotor Rehabilitation. In M. H. Thaut & D. A. Hodges (Eds.), *The Oxford Handbook of Music and the Brain* (pp. 695-714). Oxford, England: Oxford University Press.
doi: 10.1093/oxfordhb/9780198804123.013.27
- Thaut, M.H., Kenyon, G.P., Hurt, C.P., McIntosh, G.C., & Hoemberg, V. (2002a). Kinematic optimization of spatiotemporal patterns in paretic arm training with stroke patients. *Neuropsychologia*, 40, 1073-1081.
doi: 10.1016/s0028-3932(01)00141-5
- Thaut, M.H., Schicks, W., McIntosh, G.C., & Hoemberg, V. (2002b). The role of motor imagery and temporal cuing in hemiparetic arm rehabilitation. *Neurorehabilitation and Neural Repair*, 16, 115.
- Vercruyse, S., Spildooren, J., Heremans, E., Vandenbossche, J., Levin, O., Wenderoth, N., ... & Nieuwboer, A. (2012). Freezing in Parkinson's disease: a spatiotemporal motor disorder beyond gait. *Movement Disorders*, 27(2), 254-263.
doi: 10.1002/mds.24015
- Wang, T. H., Peng, Y. C., Chen, Y. L., Lu, T. W., Liao, H. F., Tang, P. F., & Shieh, J. Y. (2013). A Home-Based Program Using Patterned Sensory Enhancement Improves Resistance Exercise Effects for Children with Cerebral Palsy: A Randomized Controlled Trial. *Neurorehabilitation and neural repair*, 27(8), 684-694.
doi: 10.1177/1545968313491001
- Whitall, J., McCombe, Waller, S., Silver, K.H., & Macko, R.F. (2000). Repetitive bilateral arm training with rhythmic auditory cueing improves motor function in chronic hemiparetic stroke. *Stroke*, 31 (10), 2390-2395.
doi: 10.1161/01.str.31.10.2390
- Yasuhara, A., Sugiyama, Y. (2001). Music therapy for children with Rett Syndrome. *Brain and Development*, 23, (1), S82-84.
doi: 10.1016/s0387-7604(01)00336-9

Therapeutic Instrumental Music Performance (TIMP)

- Altenmüller E, Marco-Pallares J, Münte TF, Schneider S. (2009). Neural reorganization underlies improvement in stroke-induced motor dysfunction by music-supported therapy. *Ann N Y Acad Sci.* (1169), 395-405.

- doi: 10.1111/j.1749-6632.2009.04580.x
- Bernatzky, G., Bernatzky, P., Hesse, H.P., Staffen, W., & Ladurner, G. (2004). Stimulating music increases motor coordination in patients afflicted by Morbus Parkinson. *Neuroscience Letters*, 361, 4-8.
doi: 10.1016/j.neulet.2003.12.022
- Bukowska, A. A. (2016). Influence of neurologic music therapy to improve the activity level in a group of patients with PD. *Nordic Journal of Music Therapy*, 25, 14.
doi: 10.1080/08098131.2016.1179888
- Bukowska, A. A., Krężałek, P., Mirek, E., Bujas, P., & Marchewka, A. (2015). Neurologic music therapy training for mobility and stability rehabilitation with Parkinson's disease—A pilot study. *Frontiers in human neuroscience*, 9.
doi: 10.3389/fnhum.2015.00710
- Chadwick, D. M., & Clark, C. A. (1980). Adapting music instruments for the physically handicapped. *Music Educators Journal*, 67(3), 56-59.
doi: 10.2307/3400621
- Chong, H. J., Cho, S. R., Jeong, E., & Kim, S. J. (2013). Finger exercise with keyboard playing in adults with cerebral palsy: A preliminary study. *Journal of exercise rehabilitation*, 9(4), 420-425.
doi: 10.12965/jer.130050
- Chong, H. J., Cho, S. R., & Kim, S. J. (2014). Hand rehabilitation using MIDI keyboard playing in adolescents with brain damage: a preliminary study. *NeuroRehabilitation*, 34(1), 147-155.
doi: 10.3233/nre-131026
- Cross, P., McLellan, M., Vomberg, E., Monga, M., & Monga, T.N. (1984). Observations on the use of music in rehabilitation of stroke patients. *Physiotherapy Canada*, 36, 197-201.
link:<https://www.stroking.ca/en/publication/cross-p-mcclellan-m-vomberg-e-monga-m-monga-t-n-1984-observations-on-the-use-of-music-in-rehabilitation-of-stroke-patients-physiotherapy-canada-364-197-201/>
- Elliott, B. (1982). *Guide to the Selection of Musical Instruments with Respect to Physical Ability and Disability*. Saint Louis, MO: MMB Music, Inc.
- Fujioka T, Ween JE, Jamali S, Stuss DT, Ross B. (2012). Changes in neuromagnetic beta-band oscillation after music-supported stroke rehabilitation. *Ann N Y Acad Sci.* (1252), 294-304.
doi: 10.1111/j.1749-6632.2011.06436.x
- Grau-Sánchez, J., Amengual, J. L., Rojo, N., Veciana de las Heras, M., Montero, J., Rubio, F., ... & Rodríguez-Fornells, A. (2013). Plasticity in the sensorimotor cortex induced by Music-supported therapy in stroke patients: a TMS study. *Frontiers in human neuroscience*, 7, 494.
doi: 10.3389/fnhum.2013.00494
- Hund-Georgiadis, M., & von Cramon, D. Y. (1999). Motor-learning-related changes in piano players and non-musicians revealed by functional magnetic-resonance signals. *Experimental Brain Research*, 125(4), 417-425.
doi: 10.1007/s002210050698
- Kojovic, M., Pareés, I., Sadnicka, A., Kassavetis, P., Rubio-Agusti, I., Saifee, T. A., ... & Bhatia, K. P. (2012). The brighter side of music in dystonia. *Archives of neurology*, 69(7), 917-919.
doi: 10.1001/archneurol.2012.33
- Lim HA, Miller K, Fabian C. (2011). The effects of therapeutic instrumental music performance on endurance level, self-perceived fatigue level, and self-perceived exertion of inpatients in physical rehabilitation. *J Music Ther.* 48(2),124-148.
doi: 10.1093/jmt/48.2.124
- Pascual-Leone, A (2001). The brain that plays music and is changed by it. *Annals of the New York Academy of Sciences*, 930(1), 315-329.
doi: 10.1111/j.1749-6632.2001.tb05741.x
- Ringebach, S. D., Zimmerman, K., Chen, C. C., Mulvey, G. M., Holzapfel, S. D., Weeks, D. J., & Thaut, M. H. (2014). Adults with Down Syndrome Performed Repetitive Movements Fast with Continuous Music Cues. *Journal of Motor Learning and Development*, 2(3), 47-54.
doi: 10.1123/jmld.2014-0040
- Rojo, N., Amengual, J., Juncadella, M., Rubio, F., Camara, E., Marco-Pallares, J., ... & Altenmüller, E. (2011). Music-supported therapy induces plasticity in the sensorimotor cortex in chronic stroke: a single-case study using multimodal imaging (fMRI-TMS). *Brain Injury*, 25(7-8), 787-793.
doi: 10.3109/02699052.2011.576305
- Schneider, S., Müunte, T., Rodriguez-Fornells, A., Sailer, M., & Altenmüller, E. (2010). Music-supported training is more efficient than functional motor training for recovery of fine motor skills in stroke patients. *Music Perception: An Interdisciplinary Journal*, 27(4), 271-280.

- doi: 10.3109/02699052.2011.576305
- Street, A. J., Fachner, J., & Magee, W. L. (2019). Upper limb rehabilitation in chronic stroke using neurologic music therapy: Two contrasting case studies to inform on treatment delivery and patient suitability. *Nordic Journal of Music Therapy*, 28(5), 382-404.
doi: 10.1080/08098131.2019.1606848
- Sutton, K. (1984). The development and implementation of a music therapy physiological measures test. *Journal of Music Therapy*, 21, 160.
doi: 10.1093/jmt/21.4.160
- Thaut, C., & Stephan, K. M. (2019). Neurologic Music Therapy in Sensorimotor Rehabilitation. In M. H. Thaut & D. A. Hodges (Eds.), *The Oxford Handbook of Music and the Brain* (pp. 695-714). Oxford, England: Oxford University Press.
doi: 10.1093/oxfordhb/9780198804123.013.27
- Turova, V., Alves-Pinto, A., Ehrlich, S., Blumenstein, T., Cheng, G., & Lampe, R. (2017). Effects of short-term piano training on measures of finger tapping, somatosensory perception and motor-related brain activity in patients with cerebral palsy. *Neuropsychiatric Disease and Treatment*, 13, 2705-2718.
doi: 10.2147/ndt.s145104
- Yoo, J. (2009). The role of therapeutic instrumental music performance in hemiparetic arm rehabilitation. *Music Therapy Perspectives*, 27(1), 16-24.
doi: 10.1093/mtp/27.1.16

Speech and Language

Melodic Intonation Therapy (MIT)

- Albert, M., Sparks, R., & Helm, N. (1973). Melodic intonation therapy for aphasics. *Archives of Neurology*, 29, 130-131.
doi: 10.1001/archneur.1973.00490260074018
- Al-Janabi, S., Nickels, L. A., Sowman, P. F., Burianová, H., Merrett, D., & Thompson, B. (2014). Augmenting melodic intonation therapy with non-invasive brain stimulation to treat impaired left-hemisphere function: two case studies. *Frontiers in psychology*, 5, 37.
doi: 10.3389/fpsyg.2014.00037
- Belin, P., Van Eeckhout, P., Zilbovicius, M., Remy, P., Francois, C., Guillaume, S., Chain, F., Rancurel, G., & Sampson, Y. (1996). Recovery from nonfluent aphasia after melodic intonation therapy. *Neurology*, 47, 1504-1511.
doi: 10.1212/wnl.47.6.1504
- Berlin, C.I. (1976). On: Melodic intonation therapy for aphasia by R.W. Sparks and A.L. Holland. *Journal of Speech and Hearing Disorders*, 41, 298-300.
doi: 10.1044/jshd.4103.298
- Bonakdarpour, B., Eftekharzadeh, A., & Ashayeri, H. (2003). Melodic Intonation Therapy with Persian aphasic patients. *Aphasiology*, 17, 75-95.
doi: 10.1080/729254891
- Boucher, V., Garcia, L. J., Fleurant, J., & Paradis, J. (2001). Variable efficacy of rhythm and tone in melody-based interventions: implications for the assumption of a right-hemisphere facilitation in non-fluent aphasia. *Aphasiology*, 15(2), 131-149.
doi: 10.1080/02687040042000098
- Breier, J. I., Randle, S., Maher, L. M., & Papanicolaou, A. C. (2010). Changes in maps of language activity activation following melodic intonation therapy using magnetoencephalography: Two case studies. *Journal of Clinical And Experimental Neuropsychology*, 32(3), 309-314
doi: 10.1080/13803390903029293
- Brown, S., Martinez, M. J., & Parsons, L. M. (2006). Music and language side by side in the brain: a PET study of the generation of melodies and sentences. *European journal of neuroscience*, 23(10), 2791-2803.
doi: 10.1111/j.1460-9568.2006.04785.x
- Cadalbert, A., Landis, T., REGARD, M., & Graves, R. E. (1994). Singing with and without words: Hemispheric asymmetries in motor control. *Journal of Clinical and Experimental Neuropsychology*, 16(5), 664-670.
doi: 10.1080/01688639408402679
- Conklyn, D., Novak, E., Boissy, A., Bethoux, F., & Chemali, K. (2012). The effects of modified melodic

- intonation therapy on nonfluent aphasia: A pilot study. *Journal of Speech, Language, and Hearing Research*, 55(5), 1463-1471.
doi: 10.1044/1092-4388(2012/11-0105)
- Cortese, M. D., Riganello, F., Arcuri, F., Pignataro, L. M., & Buglione, I. (2015). Rehabilitation of aphasia: application of melodic-rhythmic therapy to Italian language. *Frontiers in human neuroscience*, 9.
doi: 10.3389/fnhum.2015.00520
- Goldfarb, R. & Bader, E. (1979). Espousing melodic intonation therapy in aphasia rehabilitation: A case study. *International Journal of Rehabilitation Research*, 2, 333-342.
doi: 10.1097/00004356-197909000-00002
- Haro-Martínez, A. M., García-Concejero, V. E., López-Ramos, A., Maté-Arribas, E., López- Táppero, J., Lubrini, G., ... & Fuentes, B. (2017). Adaptation of melodic intonation therapy to Spanish: A feasibility pilot study, *Aphasiology* 31(11), 1333–1343.
doi: 10.1080/02687038.2017.1279731
- Hébert, S., Racette, A., Gagnon, L., & Peretz, I. (2003). Revisiting the dissociation between singing and speaking in expressive aphasia. *Brain*, 126(8), 1838-1850.
doi: 10.1093/brain/awg186
- Helfrich-Miller, K.R. (1984). Melodic intonation therapy with developmentally apraxic children. *Seminars in Speech and Language*, 5, 119-126.
doi: 10.1055/s-0028-1082518
- Hough, M. (2010). Melodic intonation therapy and aphasia: Another variation on a theme. *Aphasiology*, 24(6-8), 775-786.
doi: 10.1080/02687030903501941
- Keith, R. & Aronson, A. (1975). Singing as therapy for apraxia of speech and aphasia: Report of a case. *Brain and Language*, 2, 483-488.
doi: 10.1016/s0093-934x(75)80085-x
- Krauss, T. & Galloway, H. (1982). Melodic Intonation Therapy with language delayed apraxic children. *Journal of Music Therapy*, 19 (2), 102-113.
doi: 10.1093/jmt/19.2.102
- Lee, Y. S., Thaut, C., & Santoni, C. (2019). Neurologic Music Therapy for Speech and Language Rehabilitation. In M. H. Thaut & D. A. Hodges (Eds.), *The Oxford Handbook of Music and the Brain* (pp. 715-737). Oxford, England: Oxford University Press.
doi: 10.1093/oxfordhb/9780198804123.013.28
- Lim, K. B., Kim, Y. K., Lee, H. J., Yoo, J., Hwang, J. Y., Kim, J. A., & Kim, S. K. (2013). The therapeutic effect of neurologic music therapy and speech language therapy in post-stroke aphasic patients. *Annals of rehabilitation medicine*, 37(4), 556-562.
doi: 10.5535/arm.2013.37.4.556
- Morrow-Odom, K. L., & Swann, A. B. (2013). Effectiveness of melodic intonation therapy in a case of aphasia following right hemisphere stroke. *Aphasiology*, 27(11), 1322-1338.
doi: 10.1080/02687038.2013.817522
- Naeser, M.A. & Helm-Estabrooks, N. (1985). CT scan lesion localization and response to Melodic Intonation Therapy with nonfluent aphasia cases. *Cortex*, 21, 203-223.
doi: 10.1016/s0010-9452(85)80027-7
- Natke, U., Donath, T. M., & Kalveram, K. T. (2003). Control of voice fundamental frequency in speaking versus singing. *The Journal of the Acoustical Society of America*, 113(3), 1587-1593.
doi: 10.1121/1.1543928
- Norton, A., Zipse, L., Marchina, S. & Schlaug, G. (2009). Melodic Intonation therapy: Shared insights on how it is done and why it might help. *Annals of the New York Academy of Sciences*, 1169, 431-436.
doi: 10.1111/j.1749-6632.2009.04859.x
- Overy, K., Norton, A.C., Ozdemir, E., Helm-Estabrooks, N., & Schlaug, G. (2004). Activation of the left anterior inferior frontal gyrus after melodic intonation therapy in a Broca's aphasia patient. *Proceedings Society for Neuroscience*, 595.7. [Abstract]
- Ozdemir, E., Norton, A., Schlaug, G. (2006). Shared and distinct neural correlates of singing and speaking. *Neuroimage*, 33 (2), 628-35.
doi: 10.1016/j.neuroimage.2006.07.013
- Patel, A. D., & Daniele, J. R. (2003). An empirical comparison of rhythm in language and music. *Cognition*, 87(1), B35-B45.
doi: 10.1016/s0010-0277(02)00187-7
- Patel, AD. (2005). The relationship of music to the melody of speech and to syntactic processing disorders in

- aphasia. *Annals of the New York Academy of Sciences*, 1060, 59-70.
doi: 10.1196/annals.1360.005
- Popovici, M. (1995). Melodic intonation therapy in the verbal decoding of aphasics. *Revue Roumaine de Neurologie et Psychiatrie*, 33, 57-97.
link:<https://www.semanticscholar.org/paper/Melodic-intonation-therapy-in-the-verbal-decoding-Popovici/a938db5297a973efcd3f2fb01be368196269867d>
- Racette, A., Bard, C., Peretz, I. (2006). Making non-fluent aphasics speak: sing along! *Brain*, 129, 2571-2584.
doi: 10.1093/brain/awl250
- Schlaug, G., Marchina, S., & Norton, A. (2008). From singing to speaking: Why singing may lead to recovery of expressive language function in patients with Broca's Aphasia. *Music perception*, 25, 315-323.
doi: 10.1525/mp.2008.25.4.315
- Schlaug, G., Marchina, S. & Norton, A. (2009). Evidence for plasticity in white-matter tracts of patients with chronic Broca's aphasia undergoing intense intonation-based speech therapy. *Annals of the New York Academy of Sciences*. 1169, 385-394.
doi: 10.1111/j.1749-6632.2009.04587.x
- Schlaug, G., Norton, A., Marchina, S., Zipse, L., & Wan, C. Y. (2010). From singing to speaking: facilitating recovery from nonfluent aphasia. *Future neurology*, 5(5), 657-665.
doi: 10.2217/fnl.10.44
- Seki, K., Sugishita, M. (1983). Japanese-Applied Melodic Intonation Therapy for Broca Aphasia. *Brain and Nerve (No to Shinkei)*, 35(10): 1031-7.
link:<https://www.ncbi.nlm.nih.gov/pubmed/6651979>
- Sparks, R.W., Helm, N., & Albert, M. (1974). Aphasia rehabilitation resulting from melodic intonation therapy. *Cortex*, 10, 313-316.
doi: 10.1016/s0010-9452(74)80024-9
- Sparks, R.W. & Deck, J.W. (1994). Melodic Intonation Therapy. In R. Chapey (Ed.), *Language Intervention Strategies in Adult Aphasias*. Baltimore, MD: Williams & Wilkins, 368-386.
- Sparks, R.W. & Holland, A.L. (1976). Method: Melodic intonation therapy for aphasia. *Journal of Speech and Hearing Disorders*, 41, 298-300.
- Stahl B, Kotz SA, Henseler I, Turner R, Geyer S. (2011). Rhythm in disguise: why singing may not hold the key to recovery from aphasia. *Brain*. 134(10), 3083-3093.
doi: 10.1093/brain/awr240
- Stewart, L., Walsh, V., Frith, U., & Rothwell, J. (2001). Transcranial magnetic stimulation produces speech arrest but not song arrest. *Annals of the New York Academy of Sciences*, 930, 433-435.
doi: 10.1111/j.1749-6632.2001.tb05762.x
- Thulborn, K. R., Carpenter, P. A., Just, M. A. (1999). Plasticity of language-related brain function during recovery from stroke. *Stroke*, 30 (4), 749-754.
doi: 10.1161/01.str.30.4.749
- Van der Meulen, I., van de Sandt-Koenderman, W. M. E., Heijenbrok-Kal, M. H., Visch-Brink, E. G., & Ribbers, G. M. (2014). The efficacy and timing of Melodic Intonation Therapy in subacute aphasia. *Neurorehabilitation and neural repair*, 28(6), 536-544.
doi: 10.1177/1545968313517753
- Vines, B. W., Norton, A. C., & Schlaug, G. (2011). Non-invasive brain stimulation enhances the effects of melodic intonation therapy. *The relationship between music and language*, 124.
doi: 10.3389/fpsyg.2011.00230
- Warren, J.D., Warren, J.E., Fox, N.C., & Warrington, E.K. (2003). Nothing to say, something to sing: Primary progressive dynamic aphasia. *Neurocase*, 9, 140-153.
doi: 10.1076/neur.9.2.140.15068
- Wilson, S.J. (2006). Preserved Singing in Aphasia: A Case Study of the Efficacy of Melodic Intonation Therapy. *Music Perception*, 24(1), 23-36.
doi: 10.1525/mp.2006.24.1.23
- Yamadori, A., Osumi, Y., Masuhara, S., & Okubo, M. (1977). Preservation of singing in Broca's aphasia. *Journal of Neurology, Neurosurgery and Psychiatry*, 40 (3), 221-224.
doi: 10.1136/jnnp.40.3.221
- Zipse L, Norton A, Marchina S, Schlaug G. (2012). When right is all that is left: plasticity of right-hemisphere tracts in a young aphasic patient. *Ann N Y Acad Sci*. (1252), 237-45.
doi: 10.1111/j.1749-6632.2012.06454.x

Musical Speech Stimulation (MUSTIM)

- Basso, A., Capitani, E., & Vignolo, L. A. (1979). Influence of rehabilitation on language skills in aphasic patients: a controlled study. *Archives of Neurology*, 36(4), 190-196.
doi: 10.1001/archneur.1979.00500400044005
- Brown, S., Martinez, M. J., & Parsons, L. M. (2006). Music and language side by side in the brain: a PET study of the generation of melodies and sentences. *European journal of neuroscience*, 23(10), 2791-2803.
doi: 10.1111/j.1460-9568.2006.04785.x
- Cadalbert, A., Landis, T., REGARD, M., & Graves, R. E. (1994). Singing with and without words: Hemispheric asymmetries in motor control. *Journal of Clinical and Experimental Neuropsychology*, 16(5), 664-670.
doi: 10.1080/01688639408402679
- Cortese, M. D., Riganello, F., Arcuri, F., Pignataro, L. M., & Buglione, I. (2015). Rehabilitation of aphasia: Application of melodic-rhythmic therapy to Italian language. *Frontiers in Human Neuroscience* 9, 1–8. doi: 10.3389/fnhum.2015.00520
- Ding, N., Patel, A. D., Chen, L., Butler, H., Luo, C., & Poeppel, D. (2017). Temporal modulations in speech and music. *Neuroscience & Biobehavioral Reviews* 81(B), 181–187.
doi: 10.1016/j.neubiorev.2017.02.011
- Lucia, C.M. (1987). Toward developing a model of music therapy intervention in the rehabilitation of head trauma patients. *Music Therapy Perspectives*, 62, 34-39.
doi: 10.1093/mtp/4.1.34
- Natke, U., Donath, T. M., & Kalveram, K. T. (2003). Control of voice fundamental frequency in speaking versus singing. *The Journal of the Acoustical Society of America*, 113(3), 1587-1593.
doi: 10.1121/1.1543928
- Ozdemir, E., Norton, A., Schlaug, G. (2006). Shared and distinct neural correlates of singing and speaking. *Neuroimage*, 33 (2), 628-35.
doi: 10.1016/j.neuroimage.2006.07.013
- Patel, A. D., & Daniele, J. R. (2003). An empirical comparison of rhythm in language and music. *Cognition*, 87(1), B35-B45.
doi: 10.1016/s0010-0277(02)00187-7
- Stewart, L., Walsh, V., Frith, U., & Rothwell, J. (2001). Transcranial magnetic stimulation produces speech arrest but not song arrest. *Annals of the New York Academy of Sciences*, 930, 433-435.
doi: 10.1111/j.1749-6632.2001.tb05762.x
- Straube, T., Schulz, A., Geipel, K., Mentzel, H. J., & Miltner, W. H. (2008). Dissociation between singing and speaking in expressive aphasia: The role of song familiarity. *Neuropsychologia*, 46(5), 1505-1512.
doi: 10.1016/j.neuropsychologia.2008.01.008
- Yamadori, A., Osumi, Y., Masuhara, S., & Okubo, M. (1977). Preservation of singing in Broca's aphasia. *Journal of Neurology, Neurosurgery and Psychiatry*, 40 (3), 221-224.
doi: 10.1136/jnnp.40.3.221
- Yamaguchi S, Akanuma Y, Hatayama Y, Otera M, Meguro K. (2012). Singing therapy can be effective for a patient with severe nonfluent aphasia. *Int J Rehabil Res*. 35(1),78-81.
doi: 10.1097/mrr.0b013e32835032f8

Rhythmic Speech Cueing (RSC)

- Ballard, K. J., Wambaugh, J. L., Duffy, J. R., Layfield, C., Maas, E., Mauszycki, S., & Mc Neil, M. R. (2015). Treatment for acquired apraxia of speech: A systematic review of intervention research between 2004 and 2012. *American Journal of Speech Language Pathology* 24(2), 316–337.
doi: 10.1044/2015_ajslp-14-0118
- Bang, C. (1980). A world of sound and music. *Journal of the British Association for Teachers of the Deaf*, 4, 1-10.
- Bellaire, K., Yorkston, K.M., & Beukelman, D.R. (1986). Modification of breath patterning to increase naturalness of a mildly dysarthric speaker. *Journal of Communication Disorders*, 19, 271-280.
doi: 10.1016/0021-9924(86)90033-x
- Bidelman, G. M., & Alain, C. (2015). Musical training orchestrates coordinated neuroplasticity in auditory brainstem and cortex to counteract age-related declines in categorical vowel perception. *Journal of*

- Neuroscience*, 35(3), 1240-1249.
doi: 10.1523/jneurosci.3292-14.2015
- Brendal, B. & Ziegler, W. (2008). Effectiveness of metrical pacing in the treatment of apraxia of speech. *Aphasiology*, 22 (1), 1-26.
doi: 10.1080/02687030600965464
- Caligiuri, M.P. (1989). The influence of speaking rate on articulatory hypokinesia in Parkinsonian dysarthria. *Brain and Language*, 36, 1493-1502.
doi: 10.1016/0093-934x(89)90080-1
- Cason, N., Astésano, C., & Schön, D. (2015). Bridging music and speech rhythm: Rhythmic priming and audio-motor training affect speech perception. *Acta Psychologica* 155, 43–50.
doi: 10.1016/j.actpsy.2014.12.002
- Cason, N., Hidalgo, C., Isoard, F., Roman, S., & Schön, D. (2015). Rhythmic priming enhances speech production abilities: Evidence from prelingually deaf children. *Neuropsychology*, 29(1), 102.
doi: 10.1037/neu0000115
- Chern, A., Tillmann, B., Vaughan, C., & Gordon, R. L. (2018). New evidence of a rhythmic priming effect that enhances grammaticality judgments in children. *Journal of Experimental Child Psychology* 173, 371–379.
doi: 10.1016/j.jecp.2018.04.007
- Cohen, N.S. (1988). The use of superimposed rhythm to decrease the rate of speech in a brain-damaged adolescent. *Journal of Music Therapy*, 25, 85-93.
doi: 10.1093/jmt/25.2.85
- Colcord, R.D. & Adams, M.R. (1979). Voicing duration and vocal SPL changes associated with stuttering reduction during singing. *Journal of Speech and Hearing Research*, 22 (3), 468-479.
doi: 10.1044/jshr.2203.468
- Crystal, D. 1980. *Introduction to Language Pathology*. Baltimore, MD: University Park Press.
- Darrow, A.A. & Starmer, G.J. (1986). The effect of vocal training on the intonation and rate of hearing-impaired children's speech: A pilot study. *Journal of Music Therapy*, 23, 194-201.
doi: 10.1093/jmt/23.4.194
- Dworkin, J. P., Abkarian, G. G., & Johns, D. F. (1988). Apraxia of speech: The effectiveness of a treatment regimen. *Journal of Speech and Hearing Disorders*, 53(3), 280-294.
doi: 10.1044/jshd.5303.280
- Gfeller, K., Witt, S. A., Kim, K. H., Adamek, M., & Coffman, D. (1999). Preliminary report of a computerized music training program for adult cochlear implant recipients. *Journal Academy of Rehabilitative Audiology*, 32, 11-28.
- Glover, H., Kalinowski, J., Rastatter, M., & Stuart, A. (1996). Effect of instruction to sing on stuttering frequency at normal and fast rates. *Perceptual and Motor Skills*, 83, 511-522.
doi: 10.2466/pms.1996.83.2.511
- Hammen, V.L. & Yorkston, K.M. (1996). Speech and pause characteristics following speech rate reduction in hypokinetic dysarthria. *Journal of Communication Disorders*, 29, 429-445.
doi: 10.1016/0021-9924(95)00037-2
- Hammen, V.L., Yorkston, K.M., & Minifie, F.D. (1994). Effects of temporal alterations on speech intelligibility in Parkinsonian dysarthria. *Journal of Speech and Hearing Research*, 37, 244-253.
doi: 10.1044/jshr.3702.244
- Healey, E.C., Mallard, A.R., & Adams, M.R. (1976). Factors contributing to the reduction of stuttering during singing. *Journal of Speech and Hearing Research*, 19, 475-480.
doi: 10.1044/jshr.1903.475
- Hidalgo, C., Falk, S., & Schön, D. (2017). Speak on time! Effects of a musical rhythmic training on children with hearing loss. *Hearing Research* 351, 11–18.
doi: 10.1016/j.heares.2017.05.006
- Jungblut M, Huber W, Pustelniak M, Schnitker R. (2012). The impact of rhythm complexity on brain activation during simple singing: an event-related fMRI study. *Restor Neurol Neurosci*. 30(1),39-53.
doi: 10.3233/rnn-2011-0619
- Kotz, S. A., & Gunter, T. C. (2015). Can rhythmic auditory cuing remediate language-related deficits in Parkinson's disease? *Annals of the New York Academy of Sciences*, 1337(1), 62-68.
doi: 10.1111/nyas.12657
- Kotz, S. A., & Schwartze, M. (2016). Motor timing and sequencing in speech production: A general-purpose framework. In G. Hickok & S. L. Small (Eds.), *Neurobiology of Language* (pp. 717–724). New York: Academic Press.
doi: 10.1016/b978-0-12-407794-2.00057-2

- Max, L., Yudman, Elana A. (2003). Accuracy and Variability of Isochronous Rhythmic Timing Across Motor Systems In Stuttering versus Nonstuttering Individuals. *Journal of Speech Language and Hearing Research, 46*(1): 146-63, 2003.
doi: 10.1044/1092-4388(2003/012)
- Perkins, W. H. (2001). Stuttering: a matter of bad timing. *Science, 294*(5543), 786-786.
doi: 10.1126/science.294.5543.786a
- Pilon, M.A., McIntosh, K.W., & Thaut, M.H. (1998). Auditory versus visual speech timing cues as external rate control to enhance verbal intelligibility in mixed spastic-ataxic dysarthric speakers: A pilot study. *Brain Injury, 12* (9), 793-803.
doi: 10.1080/026990598122188
- Shane, H.C. & Darley, F.L. (1978). The effect of auditory rhythmic stimulation on articulatory accuracy in apraxia of speech. *Cortex, 14* (3), 444-450.
doi: 10.1016/s0010-9452(78)80071-9
- Smith, A., & Denny, M. (1990). High-frequency oscillations as indicators of neural control mechanisms in human respiration, mastication, and speech. *Journal of Neurophysiology, 63*(4), 745-758.
doi: 10.1152/jn.1990.63.4.745
- Tamplin, J., Baker, F. A., Grocke, D., Brazzale, D. J., Pretto, J. J., Ruehland, W. R., ... & Berlowitz, D. J. (2013).
Effect of singing on respiratory function, voice, and mood after quadriplegia: a randomized controlled trial. *Archives of physical medicine and rehabilitation, 94*(3), 426-434.
doi: 10.1016/j.apmr.2012.10.006
- Thaut, M.H., McIntosh, K.H., McIntosh, G.C., & Hoemberg, V. (2001). Auditory rhythmicity enhances movement and speech motor control in patients with Parkinson's disease. *Functional Neurology, 16*, 163-172.
link: <https://www.ncbi.nlm.nih.gov/pubmed/11495422>
- Ulbrich, C. (2006). Pitch range is not pitch range. *Proc. Speech Prosody 2006*.
- Wambaugh, J.L. & Martinez, A.L. (2000). Effects of rate and rhythm control treatment on consonant production accuracy in apraxia of speech. *Aphasiology, 14*(8), 851-871.
doi: 10.1080/026870300412232
- Yorkston, K.M. & Beukelman, D.R. (1981). Ataxic dysarthria: Treatment consequences based on intelligibility and prosodic considerations. *Journal of Speech and Hearing Disorders, 46*, 398-404.
link: https://www.researchgate.net/publication/15899368_Ataxic_dysarthria_Treatment_sequences_based_on_intelligibility_and_prosodic_considerations
- Yorkston, K.M., Hammen, V.L., Beukelman, D.R., & Traynor, C.D. (1990). The effect of rate control on the intelligibility and naturalness of dysarthric speech. *Journal of Speech and Hearing Disorders, 55*, 550-560.
doi: 10.1044/jshd.5503.550

Vocal Intonation Therapy (VIT)

- Baker, F., Wigram, T., Gold, C. (2005). The effects of song-singing programme on the affective speaking intonation of people with traumatic brain injury. *Brain Injury, 19*(7), 519-28.
doi: 10.1080/02699050400005150
- Bellaire, K., Yorkston, K.M., & Beukelman, D.R. (1986). Modification of breath patterning to increase naturalness of a mildly dysarthric speaker. *Journal of Communication Disorders, 19*, 271-280.
doi: 10.1016/0021-9924(86)90033-x
- Busto-Crespo, O., Uzcanga-Lacabe, M., Abad-Marco, A., Berasategui, I., García, L., Maraví, E., ... Fernández-González, S. (2016). Longitudinal voice outcomes after voice therapy in unilateral vocal fold paralysis. *Journal of Voice 30*(6).
doi: 10.1016/j.jvoice.2015.10.018
- Darrow, A.A. & Cohen, N.S. (1991). The effect of programmed pitch practice and private instruction on the vocal reproduction accuracy of hearing-impaired children: Two case studies. *Music Therapy Perspectives, 9*, 61-65.
doi: 10.1093/mtp/9.1.61
- Di Benedetto, P., Cavazzon, M., Mondolo, F., Rugiu, G., Peratoner, A., & Biasutti, E. (2009). Voice and choral singing treatment: a new approach for speech and voice disorders in Parkinson's disease. *Eur J Phys Rehabil Med, 45*(1), 13-19.
link: <https://www.ncbi.nlm.nih.gov/pubmed/18987565>

- Gunji, A., Ishii, R., Chau, W., Kakigi, R., & Pantev, C. (2007). Rhythmic brain activities related to singing in humans. *Neuroimage*, 34(1), 426-434.
doi: 10.1016/j.neuroimage.2006.07.018
- Haneishi, E. (2001). Effects of a music therapy voice protocol on speech intelligibility, vocal acoustic measures, and mood of individuals with Parkinson's disease. *Journal of Music Therapy*, 38 (4), 273-290.
doi: 10.1093/jmt/38.4.273
- Johansson, K. et al. (2011). Effects of glossopharyngeal breathing on voice in cervical spinal cord injuries. *International Journal of Therapy and Rehabilitation*, 18(9), 501-512.
doi: 10.12968/ijtr.2011.18.9.501
- Lortie, C. L., Rivard, J., Thibeault, M., & Tremblay, P. (2017). The moderating effect of frequent singing on voice aging. *Journal of Voice* 31(1).
doi: 10.1016/j.jvoice.2016.02.015
- Natke, U., Donath, T. M., & Kalveram, K. T. (2003). Control of voice fundamental frequency in speaking versus singing. *The Journal of the Acoustical Society of America*, 113(3), 1587-1593.
doi: 10.1121/1.1543928
- Pillot, C. & Vaissiere, J. (2006). [Vocal effectiveness in speech and singing: acoustical, physiological and perceptive aspects. Applications in speech therapy.] *Revue de Laryngologie Otologie Rhinologie*, 127, 293-298.
link: <https://europepmc.org/article/med/17425003>
- Ramig, L. O., Bonitati, C. M., Lemke, J. H., & Horii, Y. (1994). Voice treatment for patients with Parkinson disease: Development of an approach and preliminary efficacy data. *Journal of Medical Speech-Language Pathology*, 2(3), 191-209.
- Russo, N. M., Nicol, T. G., Zecker, S. G., Hayes, E. A., & Kraus, N. (2005). Auditory training improves neural timing in the human brainstem. *Behavioural brain research*, 156(1), 95-103.
doi: 10.1016/j.bbr.2004.05.012
- Santoni, C., de Boer, G., Thaut, M., & Bressmann, T. (2018). Influence of altered auditory feedback on oral-nasal balance in song. *Journal of Voice*. Manuscript in print.
doi: 10.1016/j.jvoice.2018.06.014
- Tamplin, J., Baker, F. A., Grocke, D., Brazzale, D. J., Pretto, J. J., Ruehland, W. R., ... & Berlowitz, D. J. (2013).
Effect of singing on respiratory function, voice, and mood after quadriplegia: a randomized controlled trial. *Archives of physical medicine and rehabilitation*, 94(3), 426-434.
doi: 10.1016/j.apmr.2012.10.006
- Tamplin, J. (2008). A pilot study into the effect of vocal exercises and singing on dysarthric speech. *Neurorehabilitation*, 23 (3), 207-216.
doi: 10.3233/nre-2008-23302
- Tanner, M. A., Rammage, L., & Liu, L. (2016). Does singing and vocal strengthening improve vocal ability in people with Parkinson's disease? *Arts & Health* 8(3), 199-212.
doi: 10.1080/17533015.2015.1088047
- Tautscher-Basnett, A., Tomantschger, V., Keglevic, S., & Freimuller, M. (2006). Group therapy for individuals with Parkinson disease (PD) focusing on voice strengthening. *Proceedings of the 4th World Congress for Neurorehabilitation*, P1-054.

Therapeutic Singing (TS)

- Akanuma, K., Meguro, K., Satoh, M., Tashiro, M., & Itoh, M. (2016). Singing can improve speech function in aphasics associated with intact right basal ganglia and preserve right temporal glucose metabolism: Implications for singing therapy indication. *International Journal of Neuroscience* 126(1), 39-45.
doi: 10.3109/00207454.2014.992068
- Bang, C. (1980). A world of sound and music. *Journal of the British Association for Teachers of the Deaf*, 4, 1-10.
- Brown, S., Martinez, M. J., & Parsons, L. M. (2006). Music and language side by side in the brain: a PET study of the generation of melodies and sentences. *European journal of neuroscience*, 23(10), 2791-2803.
doi: 10.1111/j.1460-9568.2006.04785.x
- Darrow, A.A. & Starmer, G.J. (1986). The effect of vocal training on the intonation and rate of hearing-impaired

- children's speech: A pilot study. *Journal of Music Therapy*, 23, 194-201.
doi: 10.1093/jmt/23.4.194
- Di Benedetto, P., Cavazzon, M., Mondolo, F., Rugiu, G., Peratoner, A., & Biasutti, E. (2009). Voice and choral singing treatment: a new approach for speech and voice disorders in Parkinson's disease. *Eur J Phys Rehabil Med*, 45(1), 13-19.
link: <https://www.ncbi.nlm.nih.gov/pubmed/18987565>
- Gfeller, K., Witt, S. A., Kim, K. H., Adamek, M., & Coffman, D. (1999). Preliminary report of a computerized music training program for adult cochlear implant recipients. *Journal Academy of Rehabilitative Audiology*, 32, 11-28.
doi: 10.1179/1467010015Z.000000000269
- Glover, H., Kalinowski, J., Rastatter, M., & Stuart, A. (1996). Effect of instruction to sing on stuttering frequency at normal and fast rates. *Perceptual and Motor Skills*, 83, 511-522.
doi: 10.2466/pms.1996.83.2.511
- Gunji, A., Ishii, R., Chau, W., Kakigi, R., & Pantev, C. (2007). Rhythmic brain activities related to singing in humans. *Neuroimage*, 34(1), 426-434.
doi: 10.1016/j.neuroimage.2006.07.018
- Jackson, S.A. Treharne, D.A., & Boucher, J. (1997). Rhythm and language in children with moderate learning difficulties. *European Journal of Disorders of Communication*, 32 (1), 99-108.
doi: 10.3109/13682829709021463
- Jamaly, S., Leidag, M., Schneider, H. W., Domanski, U., Rasche, K., Schröder, M., & Nilius, G. (2017). The effect of singing therapy compared to standard physiotherapeutic lung sport in COPD. *Pneumologie* 71(S01), S1-S125.
doi: 10.1055/s-0037-1598260
- Lim, K. B., Kim, Y. K., Lee, H. J., Yoo, J., Hwang, J. Y., Kim, J. A., & Kim, S. K. (2013). The therapeutic effect of neurologic music therapy and speech language therapy in post-stroke aphasic patients. *Annals of rehabilitation medicine*, 37(4), 556-562.
doi: 10.5535/arm.2013.37.4.556
- Natke, U., Donath, T. M., & Kalveram, K. T. (2003). Control of voice fundamental frequency in speaking versus singing. *The Journal of the Acoustical Society of America*, 113(3), 1587-1593.
doi: 10.1121/1.1543928
- Racette, A., Bard, C., Peretz, I. (2006). Making non-fluent aphasics speak: sing along! *Brain*, 129, 2571-84.
doi: 10.1093/brain/awl250
- Schön, D., Boyer, M., Moreno, S., Besson, M., Peretz, I., & Kolinsky, R. (2008). Songs as an aid for language acquisition. *Cognition*, 106(2), 975-983.
doi: 10.1016/j.cognition.2007.03.005
- Shane, H.C. & Darley, F.L. (1978). The effect of auditory rhythmic stimulation on articulatory accuracy in apraxia of speech. *Cortex*, 14 (3), 444-450.
doi: 10.1016/s0010-9452(78)80071-9
- Stegemöller, E. L., Radig, H., Hibbing, P., Wingate, J., & Sapienza, C. (2017). Effects of singing on voice, respiratory control and quality of life in persons with Parkinson's disease. *Disability and Rehabilitation* 39(6), 594-600.
doi: 10.3109/09638288.2016.1152610
- Tamplin, J. (2008). A pilot study into the effect of vocal exercises and singing on dysarthric speech. *Neurorehabilitation*, 23 (3), 207-216.
doi: 10.3233/nre-2008-23302
- Tamplin, J., Baker, F. A., Grocke, D., Brazzale, D. J., Pretto, J. J., Ruehland, W. R., ... & Berlowitz, D. J. (2013). Effect of singing on respiratory function, voice, and mood after quadriplegia: a randomized controlled trial. *Archives of physical medicine and rehabilitation*, 94(3), 426-434.
doi: 10.1016/j.apmr.2012.10.006
- Wan, C. Y., Rüüber, T., Hohmann, A., & Schlaug, G. (2010). The therapeutic effects of singing in neurological disorders. *Music perception: An interdisciplinary journal*, 27(4), 287-295.
doi: 10.1525/mp.2010.27.4.287

Oral Motor and Respiratory Exercises (OMREX)

- Bang, C. (1980). A world of sound and music. *Journal of the British Association for Teachers of the Deaf*, 4, 1-10. doi: 10.1080/08098139809477935

- Darrow, A.A. & Starmer, G.J. (1986). The effect of vocal training on the intonation and rate of hearing-impaired children's speech: A pilot study. *Journal of Music Therapy*, 23, 194-201. doi: 10.1093/jmt/23.4.194
- De Swart, B. J., Willemse, S. C., Maassen, B. A. M., & Horstink, M. W. I. M. (2003). Improvement of voicing in patients with Parkinson's disease by speech therapy. *Neurology*, 60(3), 498-500. doi: 10.1212/01.WNL.0000044480.95458.56
- Haneishi, E. (2001). Effects of a music therapy voice protocol on speech intelligibility, vocal acoustic measures, and mood of individuals with Parkinson's disease. *Journal of Music Therapy*, 38 (4), 273-290. doi: 10.1093/jmt/38.4.273
- Haas, F., Distenfeld, S., & Axen, K. (1986). Effects of perceived musical rhythm on respiratory pattern. *Journal of applied physiology*, 61(3), 1185-1191. doi: 10.1152/jappl.1986.61.3.1185
- Kim, S. J., & Jo, U. (2013). Study of accent-based music speech protocol development for improving voice problems in stroke patients with mixed dysarthria. *NeuroRehabilitation*, 32(1), 185-190. doi: 10.3233/NRE-130835
- Racette, A., Bard, C., Peretz, I. (2006). Making non-fluent aphasics speak: sing along! *Brain*, 129, 2571-84. doi: 10.1093/brain/awl250
- Stegemöller, E. L., Radig, H., Hibbing, P., Wingate, J., & Sapienza, C. (2017). Effects of singing on voice, respiratory control and quality of life in persons with Parkinson's disease. *Disability and Rehabilitation* 39(6), 594-600. doi: 10.3109/09638288.2016.1152610
- Tamplin, J. (2008). A pilot study into the effect of vocal exercises and singing on dysarthric speech. *Neurorehabilitation*, 23(3), 207-216. doi: 10.3233/NRE-2008-23302
- Wambaugh, J.L. & Martinez, A.L. (2000). Effects of rate and rhythm control treatment on consonant production accuracy in apraxia of speech. *Aphasiology*, 14(8), 851-871. doi: 10.1080/026870300412232
- Wiens ME, Reimer MA, Lindsay Guyn H. (1999). Music therapy as a treatment method for improving respiratory muscle strength in patients with advanced multiple sclerosis: a pilot study. *Rehabilitation Nursing*, 24, 74-80. doi: 10.1002/j.2048-7940.1999.tb01840

Developmental Speech and Language Training through Music (DSLTM)

- Bedoin, N., Brisseau, L., Molinier, P., Roch, D., & Tillmann, B. (2016). Temporally regular musical primes facilitate subsequent syntax processing in children with specific language impairment. *Frontiers in Neuroscience*, 10. doi: 10.3389/fnins.2016.00245
- Bekius, A., Cope, T., & Grube, M. (2016). The beat to read: A cross-lingual link between rhythmic regularity perception and reading skill. *Frontiers in Human Neuroscience* 10, 425. doi:10.3389/fnhum.2016.00425
- Bolduc, J., & Lefebvre, P. (2012). Using nursery rhymes to foster phonological and musical processing skills in kindergarteners. *Creative Education*, 3(4), 495. doi: 10.4236/ce.2012.34075
- Bonacina, S., Cancer, A., Lanzi, P. L., Lorusso, M. L., & Antonietti, A. (2015). Improving reading skills in students with dyslexia: the efficacy of a sublexical training with rhythmic background. *Frontiers in psychology*, 6, 1510. doi: 10.3389/fpsyg.2015.01510
- Chen, C. C., & Miikkulainen, R. (2001). Creating melodies with evolving recurrent neural networks. In *Neural Networks, 2001. Proceedings. IJCNN'01. International Joint Conference on* (Vol. 3, pp. 2241-2246). IEEE. doi: 10.1109/IJCNN.2001.938515
- Clément, S., Planchou, C., Béland, R., Motte, J., & Samson, S. (2015). Singing abilities in children with Specific Language Impairment (SLI). *Frontiers in Psychology* 6. doi: 10.3389/fpsyg.2015.00420
- Cogo-Moreira, H., de Avila, C. R. B., Ploubidis, G. B., & de Jesus Mari, J. (2013). Effectiveness of music education for the improvement of reading skills and academic achievement in young poor readers: a pragmatic cluster-randomized, controlled clinical trial. *PloS one*, 8(3). doi:10.1371/journal.pone.0059984
- Corriveau, K.H. & Goswami, U. (2009). Rhythmic motor entrainment in children with speech and language impairments: tapping to the beat. *Cortex*, 45, 119-130. doi: 10.1016/j.cortex.2007.09.008
- Cumming, R., Wilson, A., Leong, V., Colling, L. J., & Goswami, U. (2015). Awareness of rhythm patterns in speech and music in children with specific language impairments. *Frontiers in Human Neuroscience*, 9. doi: 10.3389/fnhum.2015.00672
- Flaugnacco, E., Lopez, L., Terribili, C., Montico, M., Zoia, S., & Schön, D. (2015). Music training increases phonological awareness and reading skills in developmental dyslexia: a randomized control trial. *PloS one*, 10(9). doi:10.1371/journal.pone.0138715

- François, C., Chobert, J., Besson, M., & Schön, D. (2013). Music training for the development of speech segmentation. *Cerebral Cortex*, 23(9), 2038-2043. doi: 10.1093/cercor/bhs180
- Gfeller, K., & Baumann, A. A. (1988). Assessment procedures for music therapy with hearing impaired children: Language development. *Journal of Music Therapy*, 25(4), 192-205. doi: 10.1093/jmt/25.4.192
- Gfeller, K., & Darrow, A. A. (1987). Music as a remedial tool in the language education of hearing-impaired children. *The Arts in Psychotherapy*, 14(3), 229-235. doi: 10.1016/0197-4556(87)90006-2
- Gordon, R. L., Shivers, C. M., Wieland, E. A., Kotz, S. A., Yoder, P. J., & Devin McAuley, J. (2015). Musical rhythm discrimination explains individual differences in grammar skills in children. *Developmental Science*, 18(4), 635-644. doi: 10.1111/desc.12230
- Groß W, Linden U, Ostermann T. (2010). Effects of music therapy in the treatment of children with delayed speech development - results of a pilot study. *BMC Complementary and Alternative Medicine*. 10(1),39. doi: 10.1186/1472-6882-10-39
- Habib, M., Lardy, C., Desiles, T., Commeiras, C., Chobert, J., & Besson, M. (2016). Music and dyslexia: a new musical training method to improve reading and related disorders. *Frontiers in psychology*, 7. doi: 10.3389/fpsyg.2016.00026
- Huss M, Verney JP, Fosker T, Mead N, Goswami U. (2011). Music, rhythm, rise time perception and developmental dyslexia: perception of musical meter predicts reading and phonology. *Cortex*. 47(6),674-89. doi: 10.1016/j.cortex.2010.07.010
- Jentschke, S., Koelsch, S., Sallat, S. & Friederici, A.D. (2008). Children with specific language impairment also show impairment of music-syntactic processing. *Journal of Cognitive Neuroscience*, 20, 1940-1951. doi: 10.1162/jocn.2008.20135
- Kern, P., Wolery, M., & Aldridge, D. (2007). Use of songs to promote independence in morning greeting Routines for young children with autism. *Journal of autism and developmental disorders*, 37(7), 1264-1271. doi: 10.1007/s10803-006-0272-1
- Lai, G., Pantazatos, S. P., Schneider, H., & Hirsch, J. (2012). Neural systems for speech and song in autism. *Brain*, 135(3), 961-975. doi: 10.1093/brain/awr335
- Lim, H. A., & Draper, E. (2011). The effects of music therapy incorporated with applied behavior analysis verbal behavior approach for children with autism spectrum disorders. *Journal of music therapy*, 48(4), 532-550. doi: 10.1093/jmt/48.4.532
- Lim, H. A. (2010). Effect of “developmental speech and language training through music” on speech production in children with autism spectrum disorders. *Journal of music therapy*, 47(1), 2-26. doi.org/10.1093/jmt/47.1.2
- Linnavalli, T., Putkinen, V., Lipsanen, J., Huotilainen, M., & Tervaniemi, M. (2018). Music playschool enhances children’s linguistic skills. *Scientific Reports* 8(1), 8767. doi: 10.1038/s41598-018-27126-5
- Luria, A. R. (2012). *Higher cortical functions in man*. Springer Science & Business Media. doi: 10.1007/978-1-4684-7741-2
- Martens MA, Jungers MK, Steele AL. (2011). Effect of musical experience on verbal memory in William’s syndrome: evidence from a novel word learning task. *Neuropsychologia*. 49(11), 3093-102. doi: 10.1016/j.neuropsychologia.2011.07.016
- Milovanov, R., Huotilainen, M., Välimäki, V., Esquef, P. A., & Tervaniemi, M. (2008). Musical aptitude and second language pronunciation skills in school-aged children: neural and behavioral evidence. *Brain research*, 1194, 81-89. doi: 10.1016/j.brainres.2007.11.042
- Moore, E., Branigan, H. & Overy, K. (2017). Exploring the role of auditory-motor synchronisation in the transfer of music to language skills in dyslexia. Outstanding Poster Award talk at Neurosciences and Music VI conference.
- Moreno, S. (2009). Can music influence language and cognition? *Contemporary Music Review*, 28(3), 329-345. doi: 10.1080/07494460903404410
- Overy, K., Nicolson, R.I., Fawcett, A.J., & Clarke, E.F. (2003). Dyslexia and music: Measuring musical timing skills. *Dyslexia*, 9(1), 18-36. doi: 10.1002/dys.233
- Overy, K. (2000). Dyslexia, temporal processing and music: The potential of music as an early learning aid for dyslexic children. *Psychology of music*, 28(2), 218-229. doi: 10.1177/0305735600282010
- Saffran, J. R. (2003). Statistical language learning mechanisms and constraints. *Current directions in psychological science*, 12(4), 110-114. doi: 10.1111/1467-8721.01243
- Sandiford, G. A., Mainess, K. J., & Daher, N. S. (2013). A pilot study on the efficacy of melodic based communication therapy for eliciting speech in nonverbal children with autism. *Journal of autism and developmental disorders*, 43(6), 1298-1307. doi: 10.1007/s10803-012-1672-z
- Saylor S, Sidener TM, Reeve SA, Fetherston A, Progar PR. (2012). Effects of three types of noncontingent auditory stimulation on vocal stereotypy in children with autism. *J Appl Behav Anal*. 45(1),185-90. doi: 10.1901/jaba.2012.45-185

- Schön, D., Boyer, M., Moreno, S., Besson, M., Peretz, I., & Kolinsky, R. (2008). Songs as an aid for language acquisition. *Cognition*, 106(2), 975-983. doi: 10.1016/j.cognition.2007.03.005
- Sharda, M., Tuerk, C., Chowdhury, R. Jamey, K., Foster, N., Custo-Blanch, M., ... & Hyde, K. (2018). Music improves social communication and auditory-motor connectivity in children with autism. *Translational Psychiatry*, 8, 231. doi:10.1038/s41398-018-0287-3
- Simpson, K., Keen, D., & Lamb, J. (2013). The use of music to engage children with autism in a receptive labelling task. *Research in Autism Spectrum Disorders*, 7(12), 1489-1496. doi:10.1016/j.rasd.2013.08.013
- Slevc, L. R., Faroqi-Shah, Y., Saxena, S., & Okada, B. M. (2016). Preserved processing of musical structure in a person with agrammatic aphasia. *Neurocase* 22(6), 505-511. doi: 10.1080/13554794.2016.1177090
- Trainor, L. J., Shahin, A., & Roberts, L. E. (2003). Effects of musical training on the auditory cortex in children. *Annals of the New York Academy of Sciences*, 999(1), 506-513. doi: 10.1196/annals1284.061
- Wan, C. et al. (2011). The therapeutic effects of singing in neurological disorders. *Music Perception*, 27(4), 287-295. doi: 10.1525/mp.2010.27.4.287
- Wan, C. Y., & Schlaug, G. (2010). Neural pathways for language in autism: the potential for music-based treatments. *Future neurology*, 5(6), 797-805. doi: 10.2217/fnl.10.55
- Zuk, J., Bishop-Liebler, P., Ozernov-Palchik, O., Moore, E., Overy, K., Welch, G., & Gaab, N. (2017). Revisiting the “enigma” of musicians with dyslexia: Auditory sequencing and speech abilities. *Journal of Experimental Psychology: General*, 146(4), 495. doi: 10.1037/xge0000281

**** Cochlear implant:**

- Fu, Q.-J., Galvin, J. J., Wang, X., & Wu, J. L. (2015). Benefits of music training in Mandarin-speaking pediatric cochlear implant users. *Journal of Speech, Language, and Hearing Research* 58(1), 163-169. doi: 10.1044/2014_JSLHR-H-14-0127
- Gfeller, K. (2016). Music-based training for pediatric CI recipients: A systematic analysis of published studies. *European Annals of Otorhinolaryngology, Head and Neck Diseases*, 12th European Symposium on Pediatric Cochlear Implant (ESPCI 2015) 133(Suppl. 1), S50-S56. doi: 10.1016/j.anorl.2016.01.010
- Petersen, B., Weed, E., Sandmann, P., Brattico, E., Hansen, M., Sørensen, S. D., & Vuust, P. (2015). Brain responses to musical feature changes in adolescent cochlear implant users. *Frontiers in Human Neuroscience*, 9. doi: 10.3389/fnhum.2015.00007
- Saliba, J., Bortfeld, H., Levitin, D. J., & Oghalai, J. S. (2016). Functional near-infrared spectroscopy for neuroimaging in cochlear implant recipients. *Hearing Research*, 338(Suppl. C), 64-75. doi: 10.1016/j.heares.2016.02.005

Cognition

Musical Sensory Orientation Training (MSOT)

- Andrews, M.W. & Dowling, W.J. (1991). The development of perception of interleaved melodies and control of auditory attention. *Music Perception*, 8(4), 349-368. doi: 10.2307/40285518
- Breitling, D. et al. 1987. Auditory perception of music measured by brain electrical activity mapping. *Neuropsychologia*, 25, 765-774. doi: 10.1016/0028-3932(87)90114-X
- Castro, M., Tillmann, B., Luauté, J., Corneyllie, A., Dailier, F., André-Obadia, N., & Perrin, F. (2015). Boosting cognition with music in patients with disorders of consciousness. *Neurorehabilitation and neural repair*, 29(8), 734-742. doi: 10.1177/1545968314565464
- Cooke, M. L., Moyle, W., Shum, D. H., Harrison, S. D., & Murfield, J. E. (2010). A randomized controlled trial exploring the effect of music on agitated behaviours and anxiety in older people with dementia. *Aging and mental health*, 14(8), 905-916. doi: 10.1080/13607861003713190
- Cox, A. (2016). *Music and embodied cognition*. Bloomington, IN: Indiana University Press.
- Graham, Janet. (2004). Communication with the uncommunicative: Music therapy with pre-verbal adults. *British Journal of Learning Disabilities*, 32(1), 24-29. doi: 10.1111/j.1468-3156.2004.00247.x
- Gregory, D. (2003). Music listening for maintaining attention of older adults with cognitive impairments. *Journal of Music Therapy*, 39 (4), 244-264. doi: 10.1093/jmt/39.4.244
- Gronwall, D. (1977). Paced Auditory Serial-Addition Task: A measure of recovery from concussion. *Perceptual and Motor Skills*, 44, 367-373. doi: 10.2466/pms.1977.44.2.367
- Hammar, L. M., Götell, E., & Engström, G. (2011). Singing while caring for persons with dementia. *Arts &*

- Health*, 3(01), 39-50. doi: 10.1080/17533015.2010.481289
- Jones, M.R. (1992). Attending to musical events. In M.R. Jones & S. Holleran (Eds.), *Cognitive Bases of Musical Communication* (pp. 91-110). Washington, D.C.: American Psychological Association. doi: 10.1037/10104-006
- Korhan, E. A., Khorshid, L., & Uyar, M. (2011). The effect of music therapy on physiological signs of anxiety in patients receiving mechanical ventilatory support. *Journal of clinical nursing*, 20(7-8), 1026-1034. doi: 10.1111/j.1365-2702.2010.03434.x Ci
- Lee, H., & Noppeney, U. (2011). Long-term music training tunes how the brain temporally binds signals from multiple senses. *Proceedings of the National Academy of Sciences*, 108(51), E1441-E1450. doi: 10.1073/pnas.1115267108
- Magee WL. (2005). Music therapy with patients in low awareness states: approaches to assessment and treatment in multidisciplinary care. *Neuropsychol Rehabil.* 15(3-4), 522-36. doi: 10.1080/09602010443000461
- Magee, WL. (2007). Music as a diagnostic tool in low awareness states: Considering limbic responses. *Brain Injury*, 21, 593-599. doi: 10.1080/02699050701426907
- Noda, R., Maeda, Y., & Yoshino, A. (2004). Therapeutic time window for musicokinetic therapy in a persistent vegetative state after severe brain damage. *Brain Injury*, 18 (5), 509-515. doi: 10.1080/02699050310001645810
- Ogata, S. (1995). Human EEG responses to classical music and simulated white noise: effects of a musical loudness component on consciousness. *Perceptual and motor skills*, 80(3), 779-790. doi: 10.2466/pms.1995.80.3.779
- Okumura, Y., Asano, Y., Takenaka, S., Fukuyama, S., Yonezawa, S., Kasuya, Y., & Shinoda, J. (2014). Brain activation by music in patients in a vegetative or minimally conscious state following diffuse brain injury. *Brain injury*, 28(7), 944-950. doi: 10.3109/02699052.2014.888477
- Parenté, R., & Herrmann, D. (1996). Retraining Memory Strategies. *Topics in Language Disorders*, 17(1), 45-57.
- Park, S., Williams, R. A., & Lee, D. (2016). Effect of preferred music on agitation after traumatic brain injury. *Western journal of nursing research*, 38(4), 394-410. doi: 10.1177/0193945915593180
- Petacchi, A., Laird, A. R., Fox, P. T., & Bower, J. M. (2005). Cerebellum and auditory function: An ALE meta-analysis of functional neuroimaging studies. *Human brain mapping*, 25(1), 118-128. doi: 10.1002/hbm.20137
- Purdie, H. (1997). Music therapy in neurorehabilitation: Recent developments and new challenges. *Critical Reviews in Physical and Rehabilitation Medicine*, 9, 205-217. doi: 10.1615/CritRevPhysRehabilMed.v9.i3-4.20
- Sacks, O. (1998). Music and the Brain. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 1-18). St. Louis, MO: MMB Music, Inc.
- Sung, H. C., Lee, W. L., Li, T. L., & Watson, R. (2012). A group music intervention using percussion instruments with familiar music to reduce anxiety and agitation of institutionalized older adults with dementia. *International journal of geriatric psychiatry*, 27(6), 621-627. doi: 10.1002/gps.2761

Musical Neglect Training (MNT)

- Abiru, M., Mihara, Y., Kikuchi, Y., et al. (2007). The effects of Neurologic music therapy on hemispatial neglect in a hemiparetic stroke patient. A case study. *Neurological Medicine*, 67(1), 88-94.
- Andrews, M.W. & Dowling, W.J. (1991). The development of perception of interleaved melodies and control of auditory attention. *Music Perception*, 8(4), 349-368. doi: 10.2307/40285518
- Bernardi, N. F., Cioffi, M. C., Ronchi, R., Maravita, A., Bricolo, E., Zigiotta, L., ... & Vallar, G. (2017). Improving left spatial neglect through music scale playing. *Journal of neuropsychology*, 11, 135-158. doi: 10.1111/jnp.12078
- Bodak, R., Malhotra, P., Bernardi, N. F., Cocchini, G., & Stewart, L. (2014). Reducing chronic visuo-spatial neglect following right hemisphere stroke through instrument playing. *Frontiers in human neuroscience*, 8, 413. doi: 10.3389/fnhum.2014.00413
- Frassinetti, F., Pavani, F., & Ladavas, E. (2002). Acoustical vision of neglected stimuli: interaction among spatially converging audiovisual inputs in neglect patients. *Journal of Cognitive Neuroscience*, 14(1), 62-69. doi: 10.1162/089892902317205320
- Giacobbe, G.A. (1972). Rhythm builds order in brain-damaged children. *Music Educators Journal*, 58, 40-43.
- Guilbert, A., Clément, S., & Moroni, C. (2014). Hearing and music in unilateral spatial neglect neuro-rehabilitation. *Frontiers in psychology*, 5. doi: 10.3389/fpsyg.2014.01503

- Guilbert, A., Clement, S., & Moroni, C. (2017). A rehabilitation program based on music practice for patients with unilateral spatial neglect: A single-case study. *Neurocase* 23(1), 12–21. doi: 10.1080/13554794.2016.1265652
- Hommel, M., Peres, B., Pollak, P., Memin, B., Besson, G., Gaio, J. M., & Perret, J. (1990). Effects of passive tactile and auditory stimuli on left visual neglect. *Archives of Neurology*, 47(5), 573-576. doi: 10.1001/archneur.1990.00530050097018
- Ishihara, M., Revol, P., Jacquin-Courtois, S., Mayet, R., Rode, G., Boisson, D., ... & Rossetti, Y. (2013). Tonal cues modulate line bisection performance: preliminary evidence for a new rehabilitation prospect? *Frontiers in psychology*, 4, 704. doi: 10.3389/fpsyg.2013.00704
- Jakobson, L. S., Cuddy, L. L., & Kilgour, A. R. (2003). Time tagging: A key to musicians' superior memory. *Music Perception: An Interdisciplinary Journal*, 20(3), 307-313. doi: 10.1525/mp.2003.20.3.307
- Janata, P., Tomic, S.T., & Rakowski, S.K. (2007). Characterization of music-evoked autobiographical memories. *Memory*, 15, 845-860. doi: 10.1080/09658210701734593
- Jones, M.R. (1992). Attending to musical events. In M.R. Jones & S. Holleran (Eds.), *Cognitive Bases of Musical Communication* (pp. 91-110). Washington, D.C.: American Psychological Association. doi: 10.1037/10104-006
- Purdie, H. (1997). Music therapy in neurorehabilitation: recent developments and new challenges. *Critical Reviews™ in Physical and Rehabilitation Medicine*, 9(3-4). doi: 10.1615/CritRevPhysRehabilMed.v9.i3-4.20
- Reitan, R. M., & Wolfson, D. (1989). The Seashore Rhythm test and brain functions. *The Clinical Neuropsychologist*, 3(1), 70-78. doi: 10.1080/13854048908404078
- Robertson, I.H., Manly T., Beschin, N., Daini, R., Haeske-Dewick, H., Homberg, V., Jehkonen, M., Pizzamiglio, G., Shiel, A., Weber, E. (1997). Auditory sustained attention is a marker of unilateral spatial neglect. *Neuropsychologia*, 35(12), 1527-32.
- Soto, D., Funes, M. J., Guzmán-García, A., Warbrick, T., Rotshtein, P., & Humphreys, G. W. (2009). Pleasant music overcomes the loss of awareness in patients with visual neglect. *Proceedings of the National Academy of Sciences*, 106(14), 6011-6016. doi: 10.1073/pnas.0811681106
- Tsai, P. L., Chen, M. C., Huang, Y. T., Lin, K. C., Chen, K. L., & Hsu, Y. W. (2013). Listening to classical music ameliorates unilateral neglect after stroke. *American Journal of Occupational Therapy*, 67(3), 328-335. doi: 10.5014/ajot.2013.006312

Auditory Perception Training (APT)

- Andrews, M.W. & Dowling, W.J. (1991). The development of perception of interleaved melodies and control of auditory attention. *Music Perception*, 8(4), 349-368. doi: 10.2307/40285518
- Berger, K. A. (2013). Praxis and autism: the psychomotor regulation sensory processing dimension—a report from the field. *Frontiers in Integrative Neuroscience*, 6, 129. doi: 10.3389/fnint.2012.00129
- Bettison, S. (1996). The long-term effects of auditory training on children with autism. *Journal of Autism and Developmental Disorders*, 26(3), 361-375. doi: 10.1007/BF02172480
- Biswas, A., Hegde, S., Jhunjhunwala, K., & Pal, P. K. (2016). Two sides of the same coin: Impairment in perception of temporal components of rhythm and cognitive functions in Parkinson's disease. *Basal Ganglia* 6(1), 63–70. doi: 10.1016/j.baga.2015.12.001
- Breitling, D. et al. 1987. Auditory perception of music measured by brain electrical activity mapping. *Neuropsychologia*, 25, 765-774. doi: 10.1016/0028-3932(87)90114-X
- Brincker, M., & Torres, E. B. (2013). Noise from the periphery in autism. *Frontiers in Integrative Neuroscience*, 7, 34. doi: 10.3389/fnint.2013.00034
- DePape, A. M. R., Hall, G. B., Tillmann, B., & Trainor, L. J. (2012). Auditory processing in high-functioning adolescents with autism spectrum disorder. *PloS one*, 7(9), e44084. doi: 10.1371/journal.pone.0044084
- Fujioka, T., Ross, B., Kakigi, R., Pantey C., Trainor, L.J. (2006). One year of musical training affects development of auditory cortical-evoked fields in young children. *Brain*, 129, 2593-608. doi: 10.1093/brain/awl247
- Furukawa, Y., Uehara, K., & Furuya, S. (2017). Expertise-dependent motor somatotopy of music perception. *Neuroscience Letters* 650, 97–102. doi: 10.1016/j.neulet.2017.04.033
- Gfeller, K., Jiang, D., Oleson, J. J., Driscoll, V., Olszewski, C., Knutson, J. F., ... & Gantz, B. (2012). The

effects

- of musical and linguistic components in recognition of real-world musical excerpts by cochlear implant recipients and normal-hearing adults. *Journal of music therapy*, 49(1), 68-101. doi: 10.1093/jmt/49.1.68
- Gfeller, K., Woodworth, G., Robin, D.A., Witt, S., & Knutson, J.F. (1997). Perception of rhythmic and sequential pitch patterns by normally hearing adults and adult cochlear implant users. *Ear and Hearing*, 18, 252-260.
- Giacobbe, G.A. (1972). Rhythm builds order in brain-damaged children. *Music Educators Journal*, 58, 40-43.
- Habibi, A., Cahn, B. R., Damasio, A., & Damasio, H. (2016). Neural correlates of accelerated auditory processing in children engaged in music training. *Developmental Cognitive Neuroscienc*, 21, 1-14. doi: 10.1016/j.dcn.2016.04.003
- Habibi, A., Damasio, A., Ilari, B., Veiga, R., Joshi, A. A., Leahy, R. M., ... Damasio, H. (2017). Childhood music training induces change in micro and macroscopic brain structure: Results from a longitudinal study. *Cerebral Cortex*, 1-12. doi: 10.1093/cercor/bhx286
- Hannon, E. E., Nave-Blodgett, J. E., & Nave, K. M. (2018). The developmental origins of the perception and production of musical rhythm. *Child Development Perspectives*. doi: 10.1111/cdep.12285
- Heaton, P., Hermelin, B., & Pring, L. (1988). Autism and pitch processing: A precursor for savant musical ability? *Music Perception*, 15, 291-305. doi: 10.2307/40285769
- Hubbard, T. L. (2018). Some methodological and conceptual considerations in studies of auditory imagery. *Auditory Perception and Cognition*, 1, 6-41. doi: 10.1080/25742442.2018.1499001
- Janata, P., Tomic, S.T., & Rakowski, S.K. (2007). Characterization of music-evoked autobiographical memories. *Memory*, 15, 845-860. doi: 10.1080/09658210701734593
- Jacobson, L. S., Cuddy, L. L., & Kilgour, A. R. (2003). Time tagging: A key to musicians' superior memory. *Music Perception: An Interdisciplinary Journal*, 20(3), 307-313. doi: 10.1525/mp.2003.20.3.307
- Jung, K. H., Won, J. H., Drennan, W. R., Jameyson, E., Miyasaki, G., Norton, S. J., & Rubinstein, J. T. (2012). Psychoacoustic performance and music and speech perception in prelingually deafened children with cochlear implants. *Audiology and Neurotology*, 17(3), 189-197. doi: 10.1159/000336407
- Loui, P., Grent-'t-Jong, T., Torpey, D., Woldorff, M. (2005). Effects of attention on the neural processing of harmonic syntax in Western Music. *Cognitive Brain Research*, 25 (3), 678-87. doi: 10.1016/j.cogbrainres.2005.08.019
- Marques, C., Moreno, S., Castro, S. L., & Besson, M. (2007). Musicians detect pitch violation in a foreign language better than nonmusicians: behavioral and electrophysiological evidence. *Journal of Cognitive Neuroscience*, 19(9), 1453-1463. doi: 10.1162/jocn.2007.19.9.1453
- Mitani, C., Nakata, T., Trehub, S. E., Kanda, Y., Kumagami, H., Takasaki, K., ... & Takahashi, H. (2007). Music recognition, music listening, and word recognition by deaf children with cochlear implants. *Ear and Hearing*, 28(2), 29S-33S. doi: 10.1097/AUD.0b013e318031547a
- Morton, L.L., Kershner, J.R., & Siegel, L.S. (1990). The potential for therapeutic applications of music on problems related to memory and attention. *Journal of Music Therapy*, 26, 58-70. doi: 10.1093/jmt/27.4.195
- Parbery-Clark, A., Skoe, E., Lam, C., & Kraus, N. (2009). Musician enhancement for speech-in-noise. *Ear Hear*, 30, 653-61. doi: 10.1097/AUD.0b013e3181b412e9
- Petacchi, A., Laird, A. R., Fox, P. T., & Bower, J. M. (2005). Cerebellum and auditory function: An ALE meta-analysis of functional neuroimaging studies. *Human brain mapping*, 25(1), 118-128. doi: 10.1002/hbm.20137
- Petersen, B., Mortensen, M. V., Gjedde, A., & Vuust, P. (2009). Reestablishing speech understanding through musical ear training after cochlear implantation. *Annals of the New York Academy of Sciences*, 1169(1), 437-440. doi: 10.1111/j.1749-6632.2009.04796.x
- Purdie, H. (1997). Music therapy in neurorehabilitation: Recent developments and new challenges. *Critical Reviews in Physical and Rehabilitation Medicine*, 9, 205-217. doi: 10.1615/CritRevPhysRehabilMed.v9.i3-4.20
- Reitan, R. M., & Wolfson, D. (1989). The Seashore Rhythm test and brain functions. *The Clinical Neuropsychologist*, 3(1), 70-78.
- Robin, D.A., Tranel, D. & Damasio, H. (1990). Auditory perception of temporal and spectral events in patients with focal left and right cerebral lesions. *Brain and Language*, 39, 539-555. doi: 10.1080/13854048908404078
- Russo, N. M., Nicol, T. G., Zecker, S. G., Hayes, E. A., & Kraus, N. (2005). Auditory training improves neural

- timing in the human brainstem. *Behavioural brain research*, 156(1), 95-103.
doi: 10.1016/j.bbr.2004.05.012
- Sacks, O. (1998). Music and the Brain. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 1-18). St. Louis, MO: MMB Music, Inc.
- Torres, E. B., Brincker, M., Isenhower, R. W., Yanovich, P., Stigler, K. A., Nurnberger, J. I., ... & José, J. V. (2013). Autism: the micro-movement perspective. *Frontiers in Integrative Neuroscience*, 7, 32. doi: 10.3389/fnint.2013.00032
- Torres, E. B., Yanovich, P., & Metaxas, D. N. (2013). Give spontaneity and self-discovery a chance in ASD: spontaneous peripheral limb variability as a proxy to evoke centrally driven intentional acts. *Frontiers in Integrative Neuroscience*, 7, 46. doi: 10.3389/fnint.2013.00046
- van Besouw, R. M., Howard, D. M., & Ternström, S. (2005). Towards an understanding of speech and song perception. *Logopedics Phoniatrics Vocology*, 30(3-4), 129-135. doi: 10.1080/14015430500262160

Musical Attention Control Training (MACT)

- Andrews, M.W. & Dowling, W.J. (1991). The development of perception of interleaved melodies and control of auditory attention. *Music Perception*, 8(4), 349-368. doi: 10.2307/40285518
- Barrow, I.M, Collins, J.N., & Britt, L.D. (2006). The influence of an auditory distraction on rapid naming after a mild traumatic brain injury: a longitudinal study. *Journal of Trauma- Injury Infection & Critical Care*, 61, 1142-1149. doi: 10.1097/01.ta.0000241238.70269.c1
- Ben-Pazi, H., Gross-Tsur, V., Bergman, H., Shalev, R.S., (2003). Abnormal rhythmic motor response in children with attention-deficit-hyperactivity disorder. *Developmental Medicine & Child Neurology*, 45.743-745. doi: 10.1017/S0012162203001385
- Bonnel, A. M., Faita, F., Peretz, I., & Besson, M. (2001). Divided attention between lyrics and tunes of operatic songs: Evidence for independent processing. *Attention, Perception, & Psychophysics*, 63(7), 1201-1213.
doi: 10.3758/BF03194534
- Ceccato, E., Caneva, P., Lamonaca, D. (2006). Music therapy and cognitive rehabilitation in schizophrenic patients: A controlled study. *Nordic Journal of Music Therapy*, 15(2), 111-120.
doi: 10.1080/08098130609478158
- Cicerone, K. D., Dahlberg, C., Kalmar, K., Langenbahn, D. M., Malec, J. F., Bergquist, T. F., et al. (2000). Evidence based cognitive rehabilitation: Recommendations for clinical practice. *Archives of Physical Medicine and Rehabilitation*, 81, 1596-1615. doi: 10.1053/apmr.2000.19240
- Drake, C., Jones, M. R., & Baruch, C. (2000). The development of rhythmic attending in auditory sequences: attunement, referent period, focal attending. *Cognition*, 77(3), 251-288.
doi: 10.1016/S0010-0277(00)00106-2
- Escoffier, N., Herrmann, C. S., & Schirmer, A. (2015). Auditory rhythms entrain visual processes in the human brain: evidence from evoked oscillations and event-related potentials. *Neuroimage*, 111, 267-276.
doi: 10.1016/j.neuroimage.2015.02.024
- Gregory, D. (2003). Music listening for maintaining attention of older adults with cognitive impairments. *Journal of Music Therapy*, 39 (4), 244-264. doi: 10.1093/jmt/39.4.244
- James, T., Przynsinda, E., Sampaio, G., Woods, K. J. P., Hewett, A., Morillon, B., & Loui, P. (2017). Acoustic effects on oscillatory markers of sustained attention. Presentation at the International Conference on Auditory Cortex. Banff, Canada.
- Jones, M. R., & Ralston, J. T. (1991). Some influences of accent structure on melody recognition. *Memory & Cognition*, 19(1), 8-20. doi: 10.3758/BF03198492
- Klein, J. M., & Jones, M. R. (1996). Effects of attentional set and rhythmic complexity on attending. *Attention, Perception, & Psychophysics*, 58(1), 34-46. doi: 10.3758/BF03205473
- Knox, R., & Jutai, J. (1996). Music-based rehabilitation of attention following brain injury. *Canadian Journal of Rehabilitation*, 9, 169-181.
- Knox, R., Yokota-Adachi, H., Kershner, J., & Jutai, J. (2003). Musical attention training program and alternating attention in brain injury: An initial report. *Music Therapy Perspectives*, 21, 99-104.
doi: 10.1093/mtp/21.2.99
- Large, E. W., & Jones, M. R. (1999). The dynamics of attending: How people track time-varying events. *Psychological review*, 106(1), 119. doi: 10.1037/0033-295X.106.1.119
- Mahraun, D. (2004). The influence of music and rhythm on a sustained attention task in children with autism. (Unpublished master's thesis). Colorado State University, Fort Collins.

- Mateer, D., & New, E. (2000). 'In nomine Jesu': Robert Fayrfax and the Guild of the Holy Name in St Paul's Cathedral. *Music & letters*, 81(4), 507-519. www.jstor.org/stable/854536
- Morton, L.L., Kershner, J.R., & Siegel, L.S. (1990). The potential for therapeutic applications of music on problems related to memory and attention. *Journal of Music Therapy*, 26, 58-70. doi: 10.1093/jmt/27.4.195
- Peterson, D.A. & Thaut, M.H. (2007). Music increases frontal EEG coherence during verbal learning. *Neuroscience Letters*, 412, 217-221. doi: 10.1016/j.neulet.2006.10.057
- Purdie, H. (1997). Music therapy in neurorehabilitation: Recent developments and new challenges. *Critical Reviews in Physical and Rehabilitation Medicine*, 9, 205-217. doi: 10.1615/CritRevPhysRehabilMed.v9.i3-4.20
- Rimmele, C. T., & Hester, R. K. (1987). Cognitive rehabilitation after traumatic head injury. *Archives of Clinical Neuropsychology*, 2(4), 353-384. doi: 10.1016/0887-6177(87)90004-7
- Sacks, O. (1998). Music and the Brain. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 1-18). St. Louis, MO: MMB Music, Inc.
- Sacrey, L. A. R., Clark, C. A., & Whishaw, I. Q. (2009). Music attenuates excessive visual guidance of skilled reaching in advanced but not mild Parkinson's disease. *PLoS one*, 4(8), e6841. doi: 10.1371/journal.pone.0006841
- Särkämö, T., Tervaniemi, M., Laitinen, S., Forsblom, A., Soinila, S., Mikkonen, M., ... & Peretz, I. (2008). Music listening enhances cognitive recovery and mood after middle cerebral artery stroke. *Brain*, 131(3), 866-876. doi: 10.1093/brain/awn013
- Sohlberg, M. M., & Mateer, C. A. (1989). *Introduction to cognitive rehabilitation: Theory and practice*. Guilford Press.
- Sokolov, A. A., Miall, R. C., & Ivry, R. B. (2017). The cerebellum: Adaptive prediction for movement and cognition. *Trends in Cognitive Sciences* 21(5), 313–332. doi: 10.1016/j.tics.2017.02.005
- Strait, D. L., & Kraus, N. (2011). Can you hear me now? Musical training shapes functional brain networks for selective auditory attention and hearing speech in noise. *The relationship between music and language*, 99. doi: 10.3389/fpsyg.2011.00113
- Sussman, J.E. (2009). The effect of music on peer awareness in preschool age children with developmental disabilities. *Journal of Music Therapy*, 46, 53-68. doi: 10.1093/jmt/46.1.53
- Wolfe, D.E. & Noguchi, L.K. (2009). The use of music with young children to improve sustained attention during a vigilance task in the presence of auditory distractions. *Journal of Music Therapy*, 46, 69-83. doi: 10.1093/jmt/46.1.69

Musical Mnemonics Training (MMT)

- Agres, K., Abdallah, S., & Pearce, M. T. (2018). Information-theoretic properties of auditory sequences dynamically influence expectation and memory. *Cognitive Science* 42(1), 43–76. doi: 10.1111/cogs.12477
- Baird, A., Samson, S., Miller, L., & Chalmers, K. (2017). Does music training facilitate the mnemonic effect of song? An exploration of musicians and nonmusicians with and without Alzheimer's dementia. *Journal of Clinical and Experimental Neuropsychology*, 39(1), 9-21. doi: 10.1080/13803395.2016.1185093
- Baird, A., Umbach, H., & Thompson, W. F. (2017b). A nonmusician with severe Alzheimer's dementia learns a new song. *Neurocase*, 23(1), 36–40. doi: 10.1080/13554794.2017.1287278
- Baur, B., Uttner, I., Ilmberger, J., Fesl, G., & Mai, N. (2000). Music memory provides access to verbal knowledge in a patient with global amnesia. *Neurocase*, 6, 415-421. doi: 10.1080/13554790008402712
- Ben-Yishay, Y., Rattok, J., Ross, B., Lakin, P., Cohen, J., & Diller, L. (1980). A remedial module for the systematic amelioration of basic attentional disturbances in head trauma patients. *Working approaches to remediation of cognitive deficits in brain damaged (Rehabilitation Monograph No. 61, pp. 71-127)*. New York: New York University Medical Center, Institute of Rehabilitation Medicine.
- Ben-Yishay, Y., Piasefsky, E. B., & Rattok, J. (1987). A systematic method for ameliorating disorders in basic attention. In M. J. Meier, A. L. Benton, & L. Diller (Eds.), *Neuropsychological rehabilitation* (p. 165–181). Guilford Press.
- Berg, I. J., Koning-Haanstra, M., & Deelman, B. G. (1991). Long-term effects of memory rehabilitation: A controlled study. *Neuropsychological Rehabilitation*, 1(2), 97-111. doi: 10.1080/09602019108401384
- Bolduc, J., & Lefebvre, P. (2012). Using nursery rhymes to foster phonological and musical processing skills in kindergarteners. *Creative Education*, 3(4), 495. doi: 10.4236/ce.2012.34075
- Carruth, E. K. (1997). The effects of singing and the spaced retrieval technique on improving face-name

- recognition in nursing home residents with memory loss. *Journal of Music Therapy*, 34(3), 165-186. doi: 10.1093/jmt/34.3.165
- Cavaco, S., Feinstein, J. S., van Twillert, H., & Tranel, D. (2012). Musical memory in a patient with severe anterograde amnesia. *Journal of clinical and experimental neuropsychology*, 34(10), 1089-1100. doi: 10.1080/13803395.2012.728568
- Ceccato, E., Caneva, P., Lamonaca, D. (2006). Music therapy and cognitive rehabilitation in schizophrenic patients: A controlled study. *Nordic Journal of Music Therapy*, 15(2), 111-120. doi: 10.1080/08098130609478158
- Chan, A.S., Ho, Y.C., & Cheung, M.C. (1998). Music training improves verbal memory. *Nature*, 396, 128. doi: 10.1038/24075
- Cheung, C., Hamilton, L. S., Johnson, K., & Chang, E. F. (2016). The auditory representation of speech sounds in human motor cortex. *eLife* 5, e12577. doi: 10.7554/eLife.12577
- Cicerone, K. D., Dahlberg, C., Kalmar, K., Langenbahn, D. M., Malec, J. F., Bergquist, T. F., et al. (2000). Evidence based cognitive rehabilitation: Recommendations for clinical practice. *Archives of Physical Medicine and Rehabilitation*, 81, 1596-1615. doi: 10.1053/apmr.2000.19240
- Claussen, D. & Thaut, M.H. (1997). Music as a mnemonic device for children with learning disabilities. *Canadian Journal of Music Therapy*, 5, 55-66.
- Cook, P. R. (1999). Music, cognition, and computerized sound. The MIT Press, 167-172.
- Crystal, D. (1980). Introduction to Language Pathology. Baltimore, MD: University Park Press.
- Crystal, H. A., Grober, E., & Masur, D. A. V. I. D. (1989). Preservation of musical memory in Alzheimer's disease. *Journal of Neurology, Neurosurgery & Psychiatry*, 52(12), 1415-1416. doi: 10.1136/jnnp.52.12.1415
- Cowles, A., Beatty, W.W., Nixon, S.J., Lutz, L.J., Paulk, J., Paulk, K., & Ross, E.D. (2003). Musical skill in dementia: A violinist presumed to have Alzheimer's disease learns to play a new song. *Neurocase*, 9, 493-503. doi: 10.1076/neur.9.6.493.29378
- Cuddy, L., Duffin, J. (2005). Music, memory, and Alzheimer's disease: is music recognition spared in dementia, and how can it be assessed? *Medical Hypotheses*, 64, 229-235. doi: 10.1016/j.mehy.2004.09.005
- Deutsch, D. (1982). Organizational processes in music. In *Music, mind, and brain* (pp. 119-136). Springer US.
- El Haj, M., Clément, S., Fasotti, L., & Allain, P. (2013). Effects of music on autobiographical verbal narration in Alzheimer's disease. *Journal of Neurolinguistics*, 26(6), 691-700. doi: 10.1016/j.jneuroling.2013.06.001
- Foster, N. & Valentine, E. (2001). The effect of auditory stimulation on autobiographical recall in dementia. *Experimental Aging Research*, 27, 215-228. doi: 10.1080/036107301300208664
- Gingold, H., & Abravanel, E. (1987). Music as a mnemonic: The effects of good-and bad-music settings on verbatim recall of short passages by young children. *Psychomusicology: A Journal of Research in Music Cognition*, 7(1), 25.
- Glassman, R. B. (1999). Hypothesized neural dynamics of working memory: Several chunks might be marked simultaneously by harmonic frequencies within an octave band of brain waves. *Brain Research Bulletin*, 50(2), 77-93. doi: 10.1016/S0361-9230(99)00090-8
- Halpern, A. R., & O'connor, M. G. (2000). Implicit memory for music in Alzheimer's disease. *Neuropsychology*, 14(3), 391. doi: 10.1037/0894-4105.14.3.391
- Haslam, C. & Cook, M. (2002). Striking a chord with amnesic patient's: Evidence that song facilitates memory. *Neurocase*, 8, 453-465. doi: 10.1076/neur.8.5.453.16183
- Heaton, P., Hermelin, B., & Pring, L. (1999). Can children with autistic spectrum disorders perceive affect in music? An experimental investigation. *Psychological medicine*, 29(06), 1405-1410. doi: 10.1017/S0033291799001221
- Hermelin, B., O'Connor, N., Lee, S., & Treffert, D. (1989). Intelligence and musical improvisation. *Psychological Medicine*, 19(2), 447-457. doi: 10.1017/S0033291700012484
- Hitch, G. J. (1996). Temporal grouping effects in immediate recall: A working memory analysis. *The Quarterly Journal of Experimental Psychology: Section A*, 49(1), 116-139. doi: 10.1080/713755609
- Ho, Y. C., Cheung, M. C., & Chan, A. S. (2003). Music training improves verbal but not visual memory: cross-sectional and longitudinal explorations in children. *Neuropsychology*, 17(3), 439. doi: 10.1037/0894-4105.17.3.439
- Iwanaga, M., & Ito, T. (2002). Disturbance effect of music on processing of verbal and spatial memories. *Perceptual and motor skills*, 94(3), 1251-1258. doi: 10.2466/pms.2002.94.3c.1251
- Jakobson, L. S., Cuddy, L. L., & Kilgour, A. R. (2003). Time tagging: A key to musicians' superior memory. *Music Perception: An Interdisciplinary Journal*, 20(3), 307-313. doi: 10.1525/mp.2003.20.3.307

- Janata, P., Tillmann, B., & Bharucha, J. J. (2002). Listening to polyphonic music recruits domain-general attention and working memory circuits. *Cognitive, Affective, & Behavioral Neuroscience*, 2(2), 121-140.
doi: 10.3758/CABN.2.2.121
- Jones, M. R. (1992). Attending to musical events. In M.R. Jones and S. Holleran (Eds.), *Cognitive Bases of Musical Communication*, 91-110. Washington, DC: American Psychological Association.
doi: 10.1037/10104-006
- Kilgour, A.R., Jakobson, L.S., & Cuddy, L.L. (2000). Music training and rate of presentation as mediators of text and song recall. *Memory Cognition*, 28, 700-710. doi: 10.3758/BF03198404
- Knott, D. (2015). Musical Mnemonics Training: Proposed Mechanisms and Case Example with Acquired Brain Injury. *Music Therapy Perspectives*, miv016. doi: 10.1093/mtp/miv016
- Levitin, D. J., & Cook, P. R. (1996). Memory for musical tempo: Additional evidence that auditory memory is absolute. *Attention, Perception, & Psychophysics*, 58(6), 927-935. doi: 10.3758/BF03205494
- Ma, Y., Nagler, J., Lee, M., & Cabrera, I. (2001). Impact of music therapy on the communication skills of toddlers with pervasive developmental disorder. *Annals of the New York Academy of Sciences*, 930, 445-447. doi: 10.1111/j.1749-6632.2001.tb05766.x
- Maeller, D.H. (1996). Rehearsal strategies and verbal working memory in multiple sclerosis. Unpublished master thesis, Colorado State University.
- Mammarella, N., Fairfield, B., & Cornoldi, C. (2007). Does music enhance cognitive performance in healthy older adults? The Vivaldi effect. *Aging clinical and experimental research*, 19(5), 394.
doi: 10.1007/BF03324720
- McIntosh, G.C., Peterson, D.A., & Thaut, M.H. (2006a). Verbal learning with a musical template increases neuronal synchronization and improves verbal memory in patients with Multiple Sclerosis (MS). *Proceedings of the 4th World Congress for Neurorehabilitation*, P1-018.
- McIntosh, G.C., Peterson, D.A., & Thaut, M.H. (2006b). Verbal learning with a musical template increases neuronal synchronization and improves verbal memory in patients with multiple sclerosis. *Neurorehabilitation & Neural Repair*, 20, 129.
- Moore, K.S., Peterson, D.A., O'Shea, G., McIntosh, G.C., & Thaut, M.H. (2008). The effectiveness of music as a mnemonic device on recognition memory for people with multiple sclerosis. *Journal of Music Therapy*, 45, 307-329. doi: 10.1093/jmt/45.3.307
- Morton, L.L., Kershner, J.R., & Siegel, L.S. (1990). The potential for therapeutic applications of music on problems related to memory and attention. *Journal of Music Therapy*, 26, 58-70.
doi: 10.1093/jmt/27.4.195
- Moussard, A., Bigand, E., Belleville, S., & Peretz, I. (2014). Music as a mnemonic to learn gesture sequences in normal aging and Alzheimer's disease. *Frontiers in human neuroscience*, 8, 294.
doi: 10.3389/fnhum.2014.00294
- Palisson, J., Roussel-Baclet, C., Maillet, D., Belin, C., Ankri, J., & Narme, P. (2015). Music enhances verbal episodic memory in Alzheimer's disease. *Journal of clinical and experimental neuropsychology*, 37(5), 503-517. doi: 10.1080/13803395.2015.1026802
- Parente, R., & Anderson-Parente, J. K. (1989). Retraining memory: Theory and application. *The Journal of Head Trauma Rehabilitation*, 4(3), 55-65.
- Purdie, H. (1997). Music therapy in neurorehabilitation: Recent developments and new challenges. *Critical Reviews in Physical and Rehabilitation Medicine*, 9, 205-217.
doi: 10.1615/CritRevPhysRehabilMed.v9.i3-4.20
- Rainey, D. & Larsen, J. (2002). The effect of familiar melodies on initial learning and long-term memory for unconnected text. *Music Perception*, 20(2), 173-186. doi: 10.1525/mp.2002.20.2.173
- Sacks, O. (1998). Music and the Brain. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 1-18). St. Louis, MO: MMB Music, Inc.
- Särkämö, T., Tervaniemi, M., Laitinen, S., Forsblom, A., Soinila, S., Mikkonen, M., ... & Peretz, I. (2008). Music listening enhances cognitive recovery and mood after middle cerebral artery stroke. *Brain*, 131(3), 866-876. doi: 10.1093/brain/awn013
- Silverman, M. J. (2007). The effect of paired pitch, rhythm, and speech on working memory as measured by sequential digit recall. *Journal of Music Therapy*, 44(4), 415-427. doi: 10.1093/jmt/44.4.415
- Simmons-Stern, N. R., Budson, A. E., & Ally, B. A. (2010). Music as a memory enhancer in patients with Alzheimer's disease. *Neuropsychologia*, 48(10), 3164-3167. doi: 10.1016/j.neuropsychologia.2010.04.033
- Simmons-Stern, N. R., Deason, R. G., Brandler, B. J., Frustace, B. S., O'Connor, M. K., Ally, B. A., & Budson, A. E. (2012). Music-based memory enhancement in Alzheimer's disease: promise and limitations. *Neuropsychologia*, 50(14), 3295-3303. doi: 10.1016/j.neuropsychologia.2012.09.019

- Son, G.R., Therrien, B., & Whall, A. (2002). Implicit memory and familiarity among elders with dementia. *Journal of Nursing Scholarship*, 34, 263-267. doi: 10.1111/j.1547-5069.2002.00263.x
- Tamminen, J., Rastle, K., Darby, J., Lucas, R., & Williamson, V. J. (2017). The impact of music on learning and consolidation of novel words. *Memory* 25(1), 107–121. doi: 10.1080/09658211.2015.1130843
- Thaut, M. H., Peterson, D. A., McIntosh, G. C., & Hoemberg, V. (2014). Music mnemonics aid verbal memory and induce learning–related brain plasticity in multiple sclerosis. *Frontiers in human neuroscience*, 8, 395. doi: 10.3389/fnhum.2014.00395
- Thaut, M.H., Sena, K.M., McIntosh, G.C., & Peterson, D.A. (2007). Musical structure facilitates verbal learning in multiple sclerosis. *Musical Structure Facilitates*, 325-330. Tomaino, C.M. 1998. Music and memory: accessing residual function. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 19-27). St. Louis, MO: MMB Music, Inc. doi: 10.1525/mp.2008.25.4.325
- Tomaino, C.M. 1998. Music and memory: accessing residual function. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 19-27). St. Louis, MO: MMB Music, Inc.
- Verwey, W. B. (2001). Concatenating familiar movement sequences: The versatile cognitive processor. *Acta psychologica*, 106(1), 69-95. doi: 10.1016/S0001-6918(00)00027-5
- Vanstone, A.D., Cuddy, L.L., Duffin, J.M., & Alexander, E. (2009). Exceptional preservation of memory for tunes and lyrics: Case studies of amusia, profound deafness, and Alzheimer’s disease. *Annals of the New York Academy of Sciences*, 1169, 291-294.
- Vanstone, A.D. & Cuddy, L.L. (2010). Musical Memory in Alzheimer disease. *Aging, Neuropsychology, and Cognition*, 17, 108-128. doi: 10.1080/13825580903042676
- Vanstone, A. D., Sikka, R., Tangness, L., Sham, R., Garcia, A., & Cuddy, L. L. (2012). Episodic and semantic memory for melodies in Alzheimer's disease. *Music Perception: An Interdisciplinary Journal*, 29(5), 501-507. doi: 10.1525/mp.2012.29.5.501
- Wimporly, D., Chadwick, P., & Nash, S. (1995). Brief report: Musical interaction therapy for children with autism: An evaluative case study with two-year follow-up. *Journal of Autism and Developmental Disabilities*, 25(3), 231-248. doi: 10.1007/BF02178299
- Wolfe, D.E. & Hom, C. (1993). Use of melodies as structural prompts for learning and retention of sequential verbal information by preschool students. *Journal of Music Therapy*, 30, 100-118. doi: 10.1093/jmt/30.2.100

Associative Mood and Memory Training (AMMT)

- Alonso, I., Dellacherie, D., & Samson, S. (2015). Emotional memory for musical excerpts in young and older adults. *Frontiers in Aging Neuroscience* 23. doi:10.3389/fnagi.2015.00023
- Bailey, J. A., & Penhune, V. B. (2010). Rhythm synchronization performance and auditory working memory in early- and late-trained musicians. *Experimental Brain Research*, 204(1), 91–101. doi: 10.1007/s00221-010-2299-y
- Basaglia-Pappas, S., Laterza, M., Borg, C., Richard-Mornas, A., Favre, E., & Thomas-Antérion, C. (2013). Exploration of verbal and non-verbal semantic knowledge and autobiographical memories starting from popular songs in Alzheimer's disease. *International psychogeriatrics*, 25(05), 785-795. doi: 10.1017/S1041610212002359
- Bever, T. G. (1988). A cognitive theory of emotion and aesthetics in music. *Psychomusicology: A Journal of Research in Music Cognition*, 7(2), 165. doi: 10.1037/h0094171
- Bower, G. H. (1981). Mood and memory. *American psychologist*, 36(2), 129. doi: 10.1037/0003-066X.36.2.129
- Brown, S., Martinez, M. J., & Parsons, L. M. (2004). Passive music listening spontaneously engages limbic and paralimbic systems. *Neuroreport*, 15(13), 2033-2037.
- Carruth, E. K. (1997). The effects of singing and the spaced retrieval technique on improving face-name recognition in nursing home residents with memory loss. *Journal of Music Therapy*, 34(3), 165-186. doi: 10.1093/jmt/34.3.165
- Choppin, S., Trost, W., Dondaine, T., Millet, B., Drapier, D., Vérin, M., ... Grandjean, D. (2016). Alteration of complex negative emotions induced by music in euthymic patients with bipolar disorder. *Journal of Affective Disorders* 191, 15–23. doi: 10.1016/j.jad.2015.10.063
- Cowles, A., Beatty, W.W., Nixon, S.J., Lutz, L.J., Paulk, J., Paulk, K., & Ross, E.D. (2003). Musical skill in dementia: A violinist presumed to have Alzheimer’s disease learns to play a new song. *Neurocase*, 9, 493-503. doi: 10.1076/neur.9.6.493.29378
- Cuddy, L., Duffin, J. (2005). Music, memory, and Alzheimer’s disease: is music recognition spared in dementia, and how can it be assessed? *Medical Hypotheses*, 64, 229-235. doi: 10.1016/j.mehy.2004.09.005
- Cuddy, L. L., Duffin, J. M., Gill, S. S., Brown, C. L., Sikka, R., & Vanstone, A. D. (2012). Memory for melodies

- and lyrics in Alzheimer's disease. *Music Perception: An Interdisciplinary Journal*, 29(5), 479-491. doi: 10.1525/mp.2012.29.5.479
- Cuddy, L. L., Sikka, R., & Vanstone, A. (2015). Preservation of musical memory and engagement in healthy aging and Alzheimer's disease. *Annals of the New York Academy of Sciences*, 1337, 223–231. ISSN 0077-8923
- Demorest, S. M., Morrison, S. J., Nguyen, V. Q., & Bodnar, E. N. (2016). The influence of contextual cues on cultural bias in music memory. *Music Perception* 33(5), 590–600. doi: 10.1525/mp.2016.33.5.590
- El Haj, M. et al. (2012). Music enhances autobiographical memory in mild Alzheimer's disease. *Education Gerontology*, 38(1), 30-41. doi: 10.1080/03601277.2010.515897
- El Haj M, Fasotti L, Allain P. (2012). The involuntary nature of music-evoked autobiographical memories in Alzheimer's disease. *Conscious Cogn*. 21(1),238-46. doi: 10.1016/j.concog.2011.12.005
- Ferreri, L., & Rodriguez-Fornells, A. (2017). Music-related reward responses predict episodic memory performance. *Experimental Brain Research*, 235(12), 3721–31. doi: 10.1007/s00221-017-5095-0
- Ferreri, L., & Verga, L. (2016). Benefits of music on verbal learning and memory. *Music Perception: An Interdisciplinary Journal*, 34(2), 167–182. doi: 10.1525/mp.2016.34.2.167
- Ford, J. H., Addis, D. R., & Giovanello, K. S. (2011). Differential neural activity during search of specific and general autobiographical memories elicited by musical cues. *Neuropsychologia*, 49(9), 2514-2526. doi: 10.1016/j.neuropsychologia.2011.04.032
- Foster, N. & Valentine, E. (2001). The effect of auditory stimulation on autobiographical recall in dementia. *Experimental Aging Research*, 27, 215-228. doi: 10.1080/036107301300208664
- Frühholz, S., Trost, W., & Kotz, S. A. (2016). The sound of emotions: Towards a unifying neural network perspective of affective sound processing. *Neuroscience & Biobehavioral Reviews* 68, 96–110. doi: 10.1016/j.neubiorev.2016.05.002
- García, J. J. M., Iodice, R., Carro, J., Sánchez, J. A., Palmero, F., & Mateos, A. M. (2012). Improvement of autobiographic memory recovery by means of sad music in Alzheimer's disease type dementia. *Aging Clin. Exp. Res*, 24, 227-232. doi: 10.3275/7874
- Gingold, H., & Abravanel, E. (1987). Music as a mnemonic: The effects of good-and bad-music settings on verbatim recall of short passages by young children. *Psychomusicology: A Journal of Research in Music Cognition*, 7(1), 25.
- Gooding, L. F., Abner, E. L., Jicha, G. A., Kryscio, R. J., & Schmitt, F. A. (2014). Musical training and late-life cognition. *American journal of Alzheimer's disease and other dementias*, 29(4), 333-343. doi: 10.1177/1533317513517048
- Grady, C. L., Furey, M. L., Pietrini, P., Horwitz, B., & Rapoport, S. I. (2001). Altered brain functional connectivity and impaired short-term memory in Alzheimer's disease. *Brain*, 124(4), 739-756. doi: 10.1093/brain/124.4.739
- Halpern, A. R., & O'connor, M. G. (2000). Implicit memory for music in Alzheimer's disease. *Neuropsychology*, 14(3), 391. doi: 10.1037/0894-4105.14.3.391
- Haslam, C. & Cook, M. (2002). Striking a chord with amnesic patient's: Evidence that song facilitates memory. *Neurocase*, 8, 453-465. doi: 10.1076/neur.8.5.453.16183
- Heaton, P., Hermelin, B., & Pring, L. (1999). Can children with autistic spectrum disorders perceive affect in music? An experimental investigation. *Psychological medicine*, 29(06), 1405-1410. doi: 10.1017/S0033291799001221
- Hegde, S. (2019). Neurologic Music Therapy Targeting Cognitive and Affective Functions. In M. H. Thaut & D. A. Hodges (Eds.), *The Oxford Handbook of Music and the Brain* (pp. 738-760). Oxford, England: Oxford University Press.
- Hermelin, B., O'Connor, N., Lee, S., & Treffert, D. (1989). Intelligence and musical improvisation. *Psychological Medicine*, 19 (2), 447-457. doi: 10.1017/S0033291700012484
- Irish, M., Cunningham, C.J., Walsh, J.B., Coakley, D., Lawlor, B.A., Robertson I.H., Coen, R.F. (2006). Investigating the enhancing effect of music on autobiographical memory in mild Alzheimer's disease. *Dementia & Geriatric Cognitive Disorders*, 22(1), 108-20. doi: 10.1159/000093487
- Iwanaga, M., & Ito, T. (2002). Disturbance effect of music on processing of verbal and spatial memories. *Perceptual and motor skills*, 94(3), 1251-1258. doi: 10.2466/pms.2002.94.3c.1251
- Jakobson, L. S., Cuddy, L. L., & Kilgour, A. R. (2003). Time tagging: A key to musicians' superior memory. *Music Perception: An Interdisciplinary Journal*, 20(3), 307-313. doi: 10.1525/mp.2003.20.3.307
- Jacobsen, J. H., Stelzer, J., Fritz, T. H., Chételat, G., La Joie, R., & Turner, R. (2015). Why musical memory can be preserved in advanced Alzheimer's disease. *Brain*, 138(8), 2438-2450. doi: 10.1093/brain/awv135
- Janata, P., Tillmann, B., & Bharucha, J. J. (2002). Listening to polyphonic music recruits domain-general attention and working memory circuits. *Cognitive, Affective, & Behavioral Neuroscience*, 2(2), 121-

140.

doi: 10.3758/CABN.2.2.121

- Janata, P., Tomic, S.T., & Rakowski, S.K. (2007). Characterization of music-evoked autobiographical memories. *Memory*, 15, 845-860. doi: 10.1080/09658210701734593
- Janata P. (2009). The neural architecture of music-evoked autobiographical memories. *Cereb Cortex*. 19(11),2579-2594. doi: 10.1093/cercor/bhp008
- Jensen, U., Lewsi, B., Tranel, D., & Adolphs, R. (2004). Emotion enhances long-term declarative memory [Abstract 203.9]. *Proceedings of the Society for Neuroscience*, 203, 209.
- Levitin, D. J., & Cook, P. R. (1996). Memory for musical tempo: Additional evidence that auditory memory is absolute. *Attention, Perception, & Psychophysics*, 58(6), 927-935. doi: 10.3758/BF03205494
- Ma, Y., Nagler, J., Lee, M., & Cabrera, I. (2001). Impact of music therapy on the communication skills of toddlers with pervasive developmental disorder. *Annals of the New York Academy of Sciences*, 930, 445-447. doi: 10.1111/j.1749-6632.2001.tb05766.x
- Martin, M. A., & Metha, A. (1997). Recall of early childhood memories through musical mood induction. *The arts in Psychotherapy*, 24(5), 447-454. doi: 10.1016/S0197-4556(97)00020-8
- Markowitsch, H. J. (2000). Memory and amnesia. *Principles of cognitive and behavioral neurology*, 257-293.
- Morton, L.L., Kershner, J.R., & Siegel, L.S. (1990). The potential for therapeutic applications of music on problems related to memory and attention. *Journal of Music Therapy*, 26, 58-70. doi: 10.1093/jmt/27.4.195
- Matrone, C., & Brattico, E. (2015). The power of music on Alzheimer's disease and the need to understand the underlying molecular mechanisms. *Journal of Alzheimer's Disease and Parkinsonism* 5. doi:10.4172/2161-0460.1000196
- Peck, K. J., Girard, T. A., Russo, F. A., & Fiocco, A. J. (2016). Music and memory in Alzheimer's disease and the potential underlying mechanisms. *Journal of Alzheimer's Disease*, 51(4), 949-959. doi: 10.3233/JAD-150998
- Purdie, H. (1997). Music therapy in neurorehabilitation: Recent developments and new challenges. *Critical Reviews in Physical and Rehabilitation Medicine*, 9, 205-217. doi: 10.1615/CritRevPhysRehabilMed.v9.i3-4.20
- Rosenbaum, R.S., Furey, M.L., Horwitz, B., & Grady, C.L. (2004). Altered communication between emotion-related brain regions supports short-term memory in Alzheimer' disease. *Proceedings Society for Neuroscience*, 203.8. [Abstract]
- Sacks, O. (1998). Music and the Brain. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 1-18). St. Louis, MO: MMB Music, Inc.
- Samson, S., Dellacherie, D., & Platel, H. (2009). Emotional power of music in patients with memory disorders. *Annals of the New York Academy of Sciences*, 1169(1), 245-255.
- Särkämö, T., Tervaniemi, M., Laitinen, S., Forsblom, A., Soinila, S., Mikkonen, M., ... & Peretz, I. (2008). Music listening enhances cognitive recovery and mood after middle cerebral artery stroke. *Brain*, 131(3), 866-876. doi: 10.1093/brain/awn013
- Särkämö, T., Tervaniemi, M., & Huotilainen, M. (2013). Music perception and cognition: development, neural basis, and rehabilitative use of music. *Wiley Interdisciplinary Reviews: Cognitive Science*, 4(4), 441-451. doi: 10.1002/wcs.1237
- Talamini, F., Carretti, B., & Grassi, M. (2016). The working memory of musicians and nonmusicians. *Music Perception: An Interdisciplinary Journal* 34(2), 183-191. doi: 10.1525/mp.2016.34.2.183
- Thaut, M.H. (1988). Measuring musical responsiveness in autistic children: A comparative analysis of improvised musical tone sequences of autistic, normal, and mentally retarded individuals. *Journal of Autism and Developmental Disorders*, 18, 561-571. doi: 10.1007/BF02211874
- Tomaino, C.M. 1998. Music and memory: accessing residual function. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 19-27). St. Louis, MO: MMB Music, Inc.
- Yannou, D., Baloyiannis, S., Kostarido- Eukleides, A., Kioseoglou, G., Damianou- Marinis, E. Klonari, D., nakopoulou, E., Tzedaki, K. (2003). Music Perception in Patients with Dementia due to Alzheimer's Disease. *Proceedings of the fifth triennial ESCOM conference*, 548-551.
- Wilson, B. A., Baddeley, A., Evans, J., & Shiel, A. (1994). Errorless learning in the rehabilitation of memory impaired people. *Neuropsychological rehabilitation*, 4(3), 307-326. doi: 10.1080/09602019408401463
- Wilson, B. A., & Evans, J. J. (1996). Error-free learning in the rehabilitation of people with memory impairments. *The Journal of Head Trauma Rehabilitation*, 11(2), 54-64. doi: 10.1097/00001199-199604000-00006
- Wilson, B. A. (2004). Management and remediation of memory problems in brain-injured adults. *The essential*

handbook of memory disorders for clinicians. John Wiley & Sons, Chichester (UK), 199-226.

Wimporoy, D., Chadwick, P., & Nash, S. (1995). Brief report: Musical interaction therapy for children with autism: An evaluative case study with two-year follow-up. *Journal of Autism and Developmental Disabilities*, 25 (3), 231-248. doi: 10.1007/BF02178299

Musical Executive Function Training (MEFT)

- Balbag, M. A., Pedersen, N. L., & Gatz, M. (2014). Playing a musical instrument as a protective factor against dementia and cognitive impairment: A population-based twin study. *International Journal of Alzheimer's Disease*, 8. doi:10.1155/2014/836748
- Beaty, R. E., Benedek, M., Silvia, P. J., & Schacter, D. L. (2016). Creative cognition and brain network dynamics. *Trends in Cognitive Sciences*, 20(2), 87–95. doi: 10.1016/j.tics.2015.10.004
- Ben-Yishay, Y., Rattok, J., Ross, B., Lakin, P., Cohen, J., & Diller, L. (1980). A remedial module for the systematic amelioration of basic attentional disturbances in head trauma patients. *Working approaches to remediation of cognitive deficits in brain damaged (Rehabilitation Monograph No. 61, pp. 71-127)*. New York: New York University Medical Center, Institute of Rehabilitation Medicine.
- Ben-Yishay, Y., Piasetsky, E. B., & Rattok, J. (1987). A systematic method for ameliorating disorders in basic attention.
- Bialystok, E. & Depape, A.M. (2009). Musical expertise, bilingualism, and executive functioning. *Journal of Experimental Psychology: Human Perception & Performance*, 35, 565-574. doi: 10.1037/a0012735
- Bugos, J.A., Perlstein, W.M., McCrae, C.S., Brophy, T.S., Bedenbaugh, P.H. (2007). Individualized piano instruction enhances executive functioning and working memory in older adults. *Aging and Mental Health*, 11(4), 464-71. doi: 10.1080/13607860601086504
- Ceccato, E., Caneva, P., Lamonaca, D. (2006). Music therapy and cognitive rehabilitation in schizophrenic patients: A controlled study. *Nordic Journal of Music Therapy*, 15(2), 111-120. doi: 10.1080/08098130609478158
- Cicerone, K. D., Dahlberg, C., Kalmar, K., Langenbahn, D. M., Malec, J. F., Bergquist, T. F., et al. (2000). Evidence based cognitive rehabilitation: Recommendations for clinical practice. *Archives of Physical Medicine and Rehabilitation*, 81, 1596-1615. doi: 10.1053/apmr.2000.19240
- Evans, J. J., Emslie, H., & Wilson, B. A. (1998). External cueing systems in the rehabilitation of executive impairments of action. *Journal of the International Neuropsychological Society*, 4(4), 399-408. doi: 10.1017/S1355617798003993
- Foxx, R. M., Martella, R. C., & Marchand-Martella, N. E. (1990). The acquisition, maintenance, and generalization of problem-solving skills by closed head-injured adults. *Behavior Therapy*, 20(1), 61-76. doi: 10.1016/S0005-7894(89)80118-2
- Gardiner, J. C., & Horwitz, J. L. (2015). Neurologic music therapy and group psychotherapy for treatment of traumatic brain injury: Evaluation of a cognitive rehabilitation group. *Music Therapy Perspectives*, miu045. doi: 10.1093/mtp/miu045
- Hars, M., Herrmann, F. R., Gold, G., Rizzoli, R., & Trombetti, A. (2013). Effect of music-based multitask training on cognition and mood in older adults. *Age and ageing*, aft163. doi: 10.1093/ageing/aft163
- Heaton, P., Hermelin, B., & Pring, L. (1999). Can children with autistic spectrum disorders perceive affect in music? An experimental investigation. *Psychological medicine*, 29(06), 1405-1410. doi: 10.1017/S0033291799001221
- Hegde, S. (2014). Music-based cognitive remediation therapy for patients with traumatic brain injury. *Frontiers in neurology*, 5, 34. doi: 10.3389/fneur.2014.00034
- Hegde, S. (2019). Neurologic Music Therapy Targeting Cognitive and Affective Functions. In M. H. Thaut & D. A. Hodges (Eds.), *The Oxford Handbook of Music and the Brain* (pp. 738-760). Oxford, England: Oxford University Press. doi: 10.1093/oxfordhb/9780198804123.013.29
- Hermelin, B., O'Connor, N., Lee, S., & Treffert, D. (1989). Intelligence and musical improvisation. *Psychological Medicine*, 19(2), 447-457. doi: 10.1017/S0033291700012484
- Janus, M., Lee, Y., Moreno, S., & Bialystok, E. (2016). Effects of short-term music and second-language training on executive control. *Journal of Experimental Child Psychology*, 144, 84–97. doi: 10.1016/j.jecp.2015.11.009
- Jaschke, A. C., Honing, H., & Scherder, E. J. (2018). Longitudinal analysis of music education on executive functions in primary school children. *Frontiers in Neuroscience*, 12, 103. doi: 10.3389/fnins.2018.00103
- Ma, Y., Nagler, J., Lee, M., & Cabrera, I. (2001). Impact of music therapy on the communication skills of toddlers

- with pervasive developmental disorder. *Annals of the New York Academy of Sciences*, 930, 445-447. doi: 10.1111/j.1749-6632.2001.tb05766.x
- Raskin (2000). Executive functions. *Neuropsychological Management of Mild Traumatic Brain Injury*, 113-133. New York: Oxford University Press.
- Sacks, O. (1998). Music and the Brain. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 1-18). St. Louis, MO: MMB Music, Inc.
- Schweizer, S., Hampshire, A., Dalgleish, T., (2011). Extending Brain-Training to the Affective Domain: Increasing Cognitive and Affective Executive Control through Emotional Working Memory Training. *PLoS One*. 6(9). doi: 10.1371/journal.pone.0024372
- Slevc, L. R., Davey, N. S., Buschkuhl, M., & Jaeggi, S. M. (2016). Tuning the mind: Exploring the connections between musical ability and executive functions. *Cognition* 152, 199–211. doi: 10.1016/j.cognition.2016.03.017
- Sohlberg, M. M., Sprunk, H., & Metzelaar, K. (1988). Efficacy of an external cuing system in an individual with severe frontal lobe damage. *Cognitive Rehabilitation*, 6, 36-41
- Thaut, M.H. (1988). Measuring musical responsiveness in autistic children: A comparative analysis of improvised musical tone sequences of autistic, normal, and mentally retarded individuals. *Journal of Autism and Developmental Disorders*, 18, 561-571. doi: 10.1007/BF02211874
- Thaut, M. H., Gardiner, J. C., Holmberg, D., Horwitz, J., Kent, L., Andrews, G., ... & McIntosh, G. R. (2009). Neurologic music therapy improves executive function and emotional adjustment in traumatic brain injury rehabilitation. *Annals of the New York Academy of Sciences*, 1169(1), 406-416. doi: 10.1111/j.1749-6632.2009.04585.x
- Wimporly, D., Chadwick, P., & Nash, S. (1995). Brief report: Musical interaction therapy for children with autism: An evaluative case study with two-year follow-up. *Journal of Autism and Developmental Disabilities*, 25(3), 231-248. doi: 10.1007/BF02178299

Music Psychotherapy and Counseling (MPC)

- Ben-Yishay, Y., Rattok, J., Ross, B., Lakin, P., Cohen, J., & Diller, L. (1980). A remedial module for the systematic amelioration of basic attentional disturbances in head trauma patients. *Working approaches to remediation of cognitive deficits in brain damaged (Rehabilitation Monograph No. 61, pp. 71-127)*. New York: New York University Medical Center, Institute of Rehabilitation Medicine.
- Ben-Yishay, Y., Piasetsky, E. B., & Rattok, J. (1987). A systematic method for ameliorating disorders in basic Attention. *Neuropsychological Rehabilitation*. New York: Guilford.
- Bever, T. G. (1988). A cognitive theory of emotion and aesthetics in music. *Psychomusicology: A Journal of Research in Music Cognition*, 7(2), 165. doi: 10.1037/h0094171
- Biasutti, M. (2015). Pedagogical applications of the cognitive research on music improvisation. *Frontiers in Psychology* 6. doi: 10.3389/fpsyg.2015.00614
- Boso, M., Emanuele, E., Minazzi, V., Abbamonte, M., & Politi, P. (2007). Effect of long-term interactive music therapy on behavior profile and musical skills in young adults with severe autism. *The Journal of Alternative and Complimentary Medicine*, 13 (7), 709-712. doi: 10.1089/acm.2006.6334
- Blood, A. J., Zatorre, R. J., Bermudez, P., & Evans, A. C. (1999). Emotional responses to pleasant and unpleasant music correlate with activity in paralimbic brain regions. *Nature neuroscience*, 2(4), 382-387. doi: 10.1038/7299
- Brown, S., Martinez, M. J., & Parsons, L. M. (2004). Passive music listening spontaneously engages limbic and paralimbic systems. *Neuroreport*, 15(13), 2033-2037.
- Carr, C., d'Ardenne, P., Sloboda, A., Scott, C., Wang, D., & Priebe, S. (2012). Group music therapy for patients with persistent post-traumatic stress disorder—an exploratory randomized controlled trial with mixed methods evaluation. *Psychology and Psychotherapy: Theory, Research and Practice*, 85(2), 179-202. doi: 10.1111/j.2044-8341.2011.02026.x
- Chu, H., Yang, C. Y., Lin, Y., Ou, K. L., Lee, T. Y., O'Brien, A. P., & Chou, K. R. (2014). The impact of group music therapy on depression and cognition in elderly persons with dementia: a randomized controlled study. *Biological research for Nursing*, 16(2), 209-217. doi: 10.1177/1099800413485410
- Dellacherie D, Bigand E, Molin P, Baulac M, Samson S. (2011). Multidimensional scaling of emotional responses to music in patients with temporal lobe resection. *Cortex*. 47(9),1107-15. doi: 10.1016/j.cortex.2011.05.007
- Erkkilä, J., Gold, C., Fachner, J., Ala-Ruona, E., Punkanen, M., & Vanhala, M. (2008). The effect of improvisational music therapy on the treatment of depression: protocol for a randomised controlled

- trial. *BMC psychiatry*, 8(1), 50. doi: 10.1186/1471-244X-8-50
- Egermann, H., Fernando, N., Chuen, L., & McAdams, S. (2015). Music induces universal emotion-related psychophysiological responses: Comparing Canadian listeners to Congolese Pygmies. *Frontiers in Psychology* 5, 1341. doi: 10.3389/fpsyg.2014.01341
- Fancourt, D., Perkins, R., Ascenso, S., Carvalho, L. A., Steptoe, A., & Williamon, A. (2016). Effects of group drumming interventions on anxiety, depression, social resilience and inflammatory immune response among mental health service users. *PLoS ONE* 11(3), 1–16. doi: 10.1371/journal.pone.0151136
- Fancourt, D., Williamon, A., Carvalho, L. A., Steptoe, A., Dow, R., & Lewis, I. (2016). Singing modulates mood, stress, cortisol, cytokine and neuropeptide activity in cancer patients and carers. *Ecancermedicalscience* 10, 1–13. doi: 10.3332/ecancer.2016.631
- Finnigan, E., & Starr, E. (2010). Increasing social responsiveness in a child with autism: A comparison of music and non-music interventions. *Autism*, 14(4), 321-348. doi: 10.1177/1362361309357747
- Fischer-Terworth, C., & Probst, P. (2011). Evaluation of a TEACCH-and music therapy-based psychological intervention in mild to moderate dementia: A controlled trial. *GeroPsych: The Journal of Gerontopsychology and Geriatric Psychiatry*, 24(2), 93. doi: 10.1024/1662-9647/a000037
- Forsblom, A., Laitinen, S., Sarkamo, T. & Tervaniemi, M. (2009). Therapeutic role of music listening in stroke rehabilitation. *Annals of the New York Academy of Sciences*, 1169, 426-430. doi: 10.1111/j.1749-6632.2009.04776.x
- Gardiner, J. C., & Horwitz, J. L. (2015). Neurologic music therapy and group psychotherapy for treatment of traumatic brain injury: Evaluation of a cognitive rehabilitation group. *Music Therapy Perspectives*, 33(2), 193–201. doi: 10.1093/mtp/miu045
- Giles, G. M., & Clark-Wilson, J. (1993). *Brain injury rehabilitation: A neurobehavioural approach*. Chapman and Hall, London.
- Goerlich K.S, Witteman J, Aleman A, Martens S. (2011). speech in alexithymia, an ERP study. *PLoS One*. 6(5),e19501. doi: 10.1371/journal.pone.0019501
- Gooding, L. F., Abner, E. L., Jicha, G. A., Kryscio, R. J., & Schmitt, F. A. (2014). Musical training and late-life cognition. *American journal of Alzheimer's disease and other dementias*, 29(4), 333-343. doi: 10.1177/1533317513517048
- Gordon, W. A., & Hibbard, M. R. (1992). Critical issues in cognitive remediation. *Neuropsychology*, 6(4), 361. doi: 10.1037/0894-4105.6.4.361
- Guetin, S., Soua, B., Voiriot, G., Picot, M.C., & Herisson, C. (2009). The effect of music therapy on mood and anxiety-depression: an observational study in institutionalized patients with traumatic brain injury. *Annals of Physical & Rehabilitation Medicine*, 52, 30-40. doi: 10.1016/j.annrmp.2008.08.009
- Guzik, J. (1987). Group treatment approaches to cognitive and social needs. *Brain Injury: Cognition and Prevocational Approaches to Rehabilitation*. New York: Tiresias Press.
- Hars, M., Herrmann, F. R., Gold, G., Rizzoli, R., & Trombetti, A. (2013). Effect of music-based multitask training on cognition and mood in older adults. *Age and ageing*, aft163. doi: 10.1093/ageing/aft163
- Heaton, P., Hermelin, B., & Pring, L. (1988). Autism and pitch processing: A precursor for savant musical ability? *Music Perception*, 15, 291-305. doi: 10.2307/40285769
- Hermelin, B., O'Connor, N., Lee, S., & Treffert, D. (1989). Intelligence and musical improvisation. *Psychological Medicine*, 19(2), 447-457. doi: 10.1017/S0033291700012484
- Kay, T. (1993). Neuropsychological treatment of mild traumatic brain injury. *The Journal of Head Trauma Rehabilitation*, 8(3), 74-85. doi: 10.1097/00001199-199309000-00009
- Kantrowitz, J. T., Hoptman, M. J., Leitman, D. I., Moreno-Ortega, M., Lehrfeld, J. M., Dias, E., ... & Javitt, D. C. (2015). Neural substrates of auditory emotion recognition deficits in schizophrenia. *Journal of Neuroscience*, 35(44), 14909-14921. doi: 10.1523/JNEUROSCI.4603-14.2015
- Karageorghis, C. I., Bruce, A. C., Pottratz, S. T., Stevens, R. C., Bigliassi, M., & Hamer, M. (2017). Psychological and psychophysiological effects of recuperative music postexercise. *Medicine & Science in Sports & Exercise* 50(4), 739–746. doi: 10.1249/MSS.0000000000001497
- Katagiri, J. (2009). The effect of background music and song texts on the emotional understanding of children with autism. *Journal of Music Therapy*, 46(1), 15-31. doi: 10.1093/jmt/46.1.15
- Kern, P., Wolery, M., & Aldridge, D. (2007). Use of songs to promote independence in morning greeting routines for young children with autism. *Journal of autism and developmental disorders*, 37(7), 1264-1271. doi: 10.1007/s10803-006-0272-1
- Lepping, R. J., Atchley, R. A., Chrysiou, E., Martin, L. E., Clair, A. A., Ingram, R. E., ... & Savage, C. R. (2016). Neural processing of emotional musical and nonmusical stimuli in depression. *PLoS one*, 11(6), e0156859.
- Linnemann, A., Ditzen, B., Strahler, J., Doerr, J. M., & Nater, U. M. (2015). Music listening as a means of stress reduction in daily life. *Psychoneuroendocrinology* 60, 82–90. doi:

- 10.1016/j.psyneuen.2015.06.008
- Linnemann, A., Strahler, J., & Nater, U. M. (2016). The stress-reducing effect of music listening varies depending on the social context. *Psychoneuroendocrinology* 72, 97–105.
doi: 10.1016/j.psyneuen.2016.06.003
- Ma, Y., Nagler, J., Lee, M., & Cabrera, I. (2001). Impact of music therapy on the communication skills of toddlers with pervasive developmental disorder. *Annals of the New York Academy of Sciences*, 930, 445-447.
doi: 10.1111/j.1749-6632.2001.tb05766.x
- Maulina, T., Djustiana, N., & Shahib, M. N. (2017). The effect of music intervention on dental anxiety during dental extraction procedure. *The Open Dentistry Journal* 11(1), 565–572.
doi: 10.2174/1874210601711010565
- Mejía-Rubalcava, C., Alanís-Tavira, J., Mendieta-Zerón, H., & Sánchez-Pérez, L. (2015). Changes induced by music therapy to physiologic parameters in patients with dental anxiety. *Complementary Therapies in Clinical Practice* 21(4), 282–286. doi: 10.1016/j.ctcp.2015.10.005
- Millard, K.A.O. & Smith, J.A. (1989). The influence of group singing therapy on the behavior of Alzheimer's disease patients. *Journal of Music Therapy*, 26, 58-70. doi: 10.1093/jmt/26.2.58
- Narme, P., Tonini, A., Khatir, F., Schiaratura, L., Clément, S., Samson, S., (2012). Non-pharmacological treatment for Alzheimer's disease: comparison between musical and non-musical interventions. *Gériatrie et Psychologie Neuropsychiatrie du Vieillessement*. 10(2), 215-224. doi: 10.1684/pnv.2012.0343
- Nayak, S. et al. (2000). Effect of music therapy on mood and social interaction among individuals with acute traumatic brain injury and stroke. *Rehabilitation Psychology*, 45(3), 274-283.
doi: 10.1037/0090-5550.45.3.274
- Ozdemir, L. & Akdemir, N. (2009). Effects of multisensory stimulation on cognition, depression and anxiety levels of mildly-affected Alzheimer's patients. *Journal of the Neurological Sciences*. 283, 211-213.
doi: 10.1016/j.jns.2009.02.367
- Pearce, E., Launay, J., & Dunbar, R. I. M. (2015). The ice-breaker effect: Singing mediates fast social bonding. *Royal Society Open Science* 2, 150221. doi: 10.1098/rsos.150221
- Peng, S. M., Koo, M., & Kuo, J. C. (2010). Effect of group music activity as an adjunctive therapy on psychotic symptoms in patients with acute schizophrenia. *Archives of psychiatric nursing*, 24(6), 429-434.
doi: 10.1016/j.apnu.2010.04.001
- Prigatano, G. P. (1999). *Principles of neuropsychological rehabilitation*. Oxford University Press.
- Purdie, H. (1997). Music therapy in neurorehabilitation: Recent developments and new challenges. *Critical Reviews in Physical and Rehabilitation Medicine*, 9, 205-217.
doi: 10.1615/CritRevPhysRehabilMed.v9.i3-4.20
- Raskin, S. A., & Mateer, C. A. (1999). *Neuropsychological management of mild traumatic brain injury*. Oxford University Press.
- Reker, G. T., Birren, J., & Svensson, C. (2013). Restoring, maintaining, and enhancing personal meaning in life through autobiographical methods. *The human quest for meaning*.
- Sacks, O. (1998). Music and the Brain. In C.M. Tomaino (Ed.), *Clinical Applications of Music in Neurologic Rehabilitation* (pp. 1-18). St. Louis, MO: MMB Music, Inc.
- Särkämö, T., Tervaniemi, M., Laitinen, S., Forsblom, A., Soynila, S., Mikkonen, M., ... & Peretz, I. (2008). Music listening enhances cognitive recovery and mood after middle cerebral artery stroke. *Brain*, 131(3), 866-876. doi: 10.1093/brain/awn013
- Särkämö, T., Tervaniemi, M., & Huotilainen, M. (2013). Music perception and cognition: development, neural basis, and rehabilitative use of music. *Wiley Interdisciplinary Reviews: Cognitive Science*, 4(4), 441-451.
doi: 10.1002/wcs.1237
- Sauvé, S., Sayad, A., Dean, R. T., & Pearce, M. T. (2017). Effects of pitch and timing expectancy on musical emotion. arXiv Preprint, 1708.03687. doi: 10.1037/pmu0000203
- Seinfeld, S., Figueroa, H., Ortiz-Gil, J., & Sanchez-Vives, M. V. (2013). Effects of music learning and piano practice on cognitive function, mood and quality of life in older adults. *Frontiers in psychology*, 4, 810.
doi: 10.3389/fpsyg.2013.00810
- Sohlberg, M.M. (2000). Psychotherapy approaches, In S.A. Raskin and C.A. Mateer (Eds), *Neuropsychological management of mild traumatic brain injury*. Oxford University Press.
- Suzuki, M., Kanamori, M., Watanabe, M., Nagasawa, S., Kojima, E., Ooshiro, H., Nakahara, D. (2004). Behavioral and endocrinological evaluation of music therapy for elderly patients with dementia. *Nursing & Health Sciences*, 6(1), 11-8. doi: 10.1111/j.1442-2018.2003.00168.x
- Thaut, M.H. (1988). Measuring musical responsiveness in autistic children: A comparative analysis of

- improvised musical tone sequences of autistic, normal, and mentally retarded individuals. *Journal of Autism and Developmental Disorders*, 18, 561-571. doi: 10.1007/BF02211874
- Ulrich G, Houtmans T, Gold C. (2007). The additional therapeutic effect of group music therapy for schizophrenic patients: a randomized study. *Acta Psychiatr Scand.* 116(5),362-70. doi: 10.1111/j.1600-0447.2007.01073.x
- Unkefer, R.F. & Thaut, M.H. (Eds.). (2002). *Music Therapy in the Treatment of Adults with Mental Disorders* (2nd ed.). St. Louis, MO: MMB Music, Inc.
- Wheeler, B. et al. (2003). Effects of number of sessions and group or individual music therapy on the mood and behavior of people who have had strokes or traumatic brain injuries. *Nordic Journal of Music Therapy*, 12(2), 139-151. doi: 10.1080/08098130309478084
- Quintin, E. M., Bhatara, A., Poissant, H., Fombonne, E., & Levitin, D. J. (2011). Emotion perception in music in high-functioning adolescents with autism spectrum disorders. *Journal of autism and developmental disorders*, 41(9), 1240-1255. doi: 10.1007/s10803-010-1146-0
- Wimpory, D., Chadwick, P., & Nash, S. (1995). Brief report: Musical interaction therapy for children with autism: An evaluative case study with two-year follow-up. *Journal of Autism and Developmental Disabilities*, 25(3), 231-248. doi: 10.1007/BF02178299